

Chiral Perturbation Theory and Lattice QCD

H. Leutwyler
University of Bern

School on Quantum Field Theories on Lattice

Saha Institute of Nuclear Physics, Kolkata
2 - 6 December, 2013

I. Standard Model at low energies

1. Interactions

Local symmetries

2. QED+QCD

Precision theory for $E \ll 100$ GeV

Qualitative difference QED \iff QCD

3. Chiral symmetry

Some of the quarks happen to be light

Approximate chiral symmetry

Spontaneous symmetry breakdown

4. Goldstone theorem

If N_f of the quark masses are put equal to zero

QCD contains $N_f^2 - 1$ Nambu-Goldstone bosons

5. Gell-Mann-Oakes-Renner relation

Quark masses break chiral symmetry

NGBs pick up mass

M_π^2 is proportional to $m_u + m_d$

II. Chiral perturbation theory

6. Group geometry

Symmetry group of the Hamiltonian G

Symmetry group of the ground state H

Nambu-Goldstone bosons live on G/H

7. Generating functional of QCD

Collects the Green functions of the theory

8. Ward identities

Symmetries of the generating functional

9. Low energy expansion

Taylor series in powers of external momenta

NGBs generate infrared singularities

10. Effective Lagrangian

Singularities due to the Nambu-Goldstone bosons can be worked out with an effective field theory.

11. Explicit construction of \mathcal{L}_{eff}

III. Illustrations

12. Some tree level calculations

Leading terms of the chiral perturbation series for the quark condensate and for M_π, F_π

13. M_π beyond tree level

Contributions to M_π at NL and NNL orders

14. F_π to one loop

Chiral logarithm in F_π , low energy theorem for scalar radius

15. Pion form factors

Charge radius of the pion, scalar radius
Dispersion relations

16. Lattice results for M_π, F_π

Determination of the low energy constants l_3, l_4 on the lattice

17. $\pi\pi$ scattering

χ P T, lattice, precision experiments

18. Conclusions for $SU(2) \times SU(2)$

19. Expansion in powers of m_s

Form of the effective Lagrangian, lattice results for the LEC

20. Zweig rule

21. Quark mass ratios

22. $\eta \rightarrow 3\pi$

23. Conclusions for $SU(3) \times SU(3)$

24. $T \neq 0$

25. Finite volume

Exercises

I. Standard Model at low energies

1. Interactions

strong weak e.m. gravity

$SU(3) \times SU(2) \times U(1) \times D$

Gravity

understood only at classical level

gravitational waves ✓

quantum theory of gravity ?

classical theory adequate for distances large compared to

$$\ell_{\text{Planck}} \equiv \sqrt{\frac{G \hbar}{c^3}} = 1.6 \cdot 10^{-35} \text{ m}$$

Units

- The constants c, \hbar, ϵ_0, k can be used to express masses, lengths, times, charges, degrees in energy units.

- mass in energy units: $m = m_{\text{SI}} c^2$

$$[m_{\text{SI}}] = 1 \text{ kg}, [m] = 1 \text{ J}$$

$$1 \text{ eV} = 1.602 \dots 10^{-19} \text{ J}$$

- length in energy units: $l = \frac{\ell_{\text{SI}}}{\hbar c}$

$$[\ell_{\text{SI}}] = 1 \text{ m}, [l] = 1 \text{ J}^{-1}$$

$$1 \text{ fm}^{-1} = 197.32 \text{ MeV}$$

- velocity becomes dimensionless: $v = \frac{v_{\text{SI}}}{c}$

- charge also dimensionless: $e = \frac{e_{\text{SI}}}{\sqrt{\epsilon_0 \hbar c}}$

⇒ Fine structure constant: $\alpha = \frac{e_{\text{SI}}^2}{4\pi\epsilon_0\hbar c} = \frac{e^2}{4\pi}$

⇒ Bohr radius: $a_{\text{Bohr}} = \frac{4\pi\epsilon_0\hbar^2}{e_{\text{SI}}^2 m_{e\text{SI}}} = \frac{4\pi}{e^2 m_e}$

- In energy units, the numerical values of the four constants are $c = \hbar = \epsilon_0 = k = 1$.

- Standard Model is a precision theory for the structure of matter
 - constituents, building blocks: u, d, e
 - held together by strong and electromagnetic interactions: gluons, photons
 - at low energies, weak interaction only generates tiny, calculable corrections

Briefly discuss the qualitative properties of the weak, electromagnetic and strong interactions at low energies

Weak interaction

- frozen at low energies

$$E \ll M_w c^2 \simeq 80 \text{ GeV}$$

- heavy quarks and leptons decay into light ones
- $s, c, b, t, \mu, \tau, \nu_e, \nu_\mu, \nu_\tau$ only contribute indirectly, via quantum fluctuations

Electromagnetic interaction

- Final form of the laws obeyed by the electromagnetic field: J. C. Maxwell 1865
Royal Society Transactions 155 (1865) 459
survived relativity and quantum theory, unharmed.
- Schrödinger equation for electrons in an electromagnetic field:

$$\frac{1}{i} \frac{\partial \psi}{\partial t} - \frac{1}{2m_e^2} (\vec{\nabla} + i e \vec{A})^2 \psi - e \varphi \psi = 0$$

contains the potentials \vec{A} , φ

- only $\vec{E} = -\vec{\nabla}\varphi - \frac{\partial \vec{A}}{\partial t}$ and $\vec{B} = \vec{\nabla} \times \vec{A}$
are of physical significance

- Fock pointed out that the Schrödinger equation is invariant under a group of local transformations: Fock 1926

$$\vec{A}' = \vec{A} + \vec{\nabla} f \quad , \quad \varphi' = \varphi - \frac{\partial f}{\partial t} \quad , \quad \psi' = e^{-ief} \psi$$

describe the same physical situation as \vec{A}, φ, ψ

- Weyl termed these **gauge transformations**
- Equivalence principle of the e.m. interaction:

$$\psi \text{ physically equivalent to } e^{-ief} \psi$$

- e^{-ief} is unitary 1×1 matrix, $e^{-ief} \in U(1)$
 $f = f(\vec{x}, t)$ space-time dependent function
- gauge invariance \iff local $U(1)$ symmetry
electromagnetic field is gauge field of $U(1)$

Weyl 1929

- $U(1)$ symmetry + renormalizability
 fully determine the e.m. interaction

Strong interaction

nuclei = p + n ~ 1930

- Nuclear forces

Stueckelberg, Yukawa ~ 1935

$$V_{e.m.} = -\frac{e^2}{4\pi r} \quad V_s = -\frac{h^2}{4\pi r} e^{-\frac{r}{r_0}}$$

$$\frac{e^2}{4\pi} \simeq \frac{1}{137}$$

$$\frac{h^2}{4\pi} \simeq 13$$

long range

short range

$$r_0 = \infty$$

$$r_0 = \frac{\hbar}{M_\pi c} = 1.4 \cdot 10^{-15} \text{ m}$$

$$M_\gamma = 0$$

$$M_\pi c^2 \simeq 140 \text{ MeV}$$

- Problem with Yukawa formula:
p and n are extended objects
diameter comparable to range of force
formula only holds for $r \gg$ diameter

- Protons, neutrons composed of quarks

$$p = uud \quad n = udd$$

- Quarks carry internal quantum number

$$u = \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix} \quad d = \begin{pmatrix} d_1 \\ d_2 \\ d_3 \end{pmatrix}$$

occur in 3 “colours”

- Strong interaction is invariant under local rotations in colour space 1973

$$u' = U \cdot u \quad d' = U \cdot d$$

$$U = \begin{pmatrix} U_{11} & U_{12} & U_{13} \\ U_{21} & U_{22} & U_{23} \\ U_{31} & U_{32} & U_{33} \end{pmatrix} \in \text{SU}(3)$$

- Can only be so if the strong interaction is also mediated by a gauge field

gauge field of SU(3) \Rightarrow strong interaction

Quantum chromodynamics

Comparison of e.m. and strong interaction

	QED	QCD
symmetry	U(1)	SU(3)
gauge field	\vec{A}, φ	gluon field
particles	photons	gluons
source	charge	colour
coupling constant	e	g

- All charged particles generate e.m. field
- All coloured particles generate gluon field
- Leptons do not interact strongly because they do not carry colour
- Equivalence principle of the strong interaction:

$$U \cdot \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix} \text{ physically equivalent to } \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix}$$

2. QED+QCD

Effective theory for $E \ll M_W c^2 \simeq 80 \text{ GeV}$

Symmetry $U(1) \times SU(3)$

Lagrangian QED+QCD

- Dynamical variables:
gauge fields for photons and gluons
Fermi fields for leptons and quarks
- Interaction fully determined by group geometry
Lagrangian contains 2 coupling constants

$$e, g$$

- Quark and lepton mass matrices can be brought to diagonal form, eigenvalues real, positive

$$m_e, m_\mu, m_\tau, m_u, m_d, m_s, m_c, m_b, m_t$$

- Transformation generates vacuum angle

$$\theta$$

- Precision theory for cold matter, atomic structure, solids, ...

Bohr radius:
$$a = \frac{4\pi}{e^2 m_e}$$

- θ breaks CP

Neutron dipole moment is very small

\Rightarrow strong upper limit, $\theta \simeq 0$


Qualitative difference between e.m. and strong interactions

- Photons do not have charge
- Gluons do have colour

$x_1 \cdot x_2 = x_2 \cdot x_1$ for $x_1, x_2 \in U(1)$ abelian

$x_1 \cdot x_2 \neq x_2 \cdot x_1$ for $x_1, x_2 \in SU(3)$

⇒ Consequence for vacuum polarization




⇒ The electromagnetic and strong interactions polarize the vacuum very differently.

Comparison with gravity


- source of gravitational field: **energy**
gravitational field does carry **energy**
- source of e.m. field: **charge**
e.m. field does not carry **charge**
- source of gluon field: **colour**
gluon field does carry **colour**

Mercury + sun



gravity

π^+ -meson



strong interaction

Perihelion shift of Mercury:

$$43'' = 50'' - 7'' \text{ per century}$$

↑

- Force between u and \bar{u} :

$$V_s = -\frac{4}{3} \frac{g^2}{4\pi r}, \quad g \rightarrow 0 \quad \text{for} \quad r \rightarrow 0$$

$$\frac{g^2}{4\pi} = \frac{6\pi}{(11N_c - 2N_f) |\ln(r \Lambda_{\text{QCD}})|}$$

$$|\ln(r \Lambda_{\text{QCD}})| \simeq 7 \quad \text{for} \quad r = \frac{\hbar}{M_Z c} \simeq 2 \cdot 10^{-18} \text{ m}$$

- Vacuum amplifies gluonic field of a bare quark
 - Field energy surrounding isolated quark = ∞
Only colour neutral states have finite energy
- \Rightarrow Confinement of colour
- \nexists analytic proof that QCD does confine colour.
Very good evidence from numerical simulations on a lattice.

QED: interaction weak at low energies

QCD: interaction strong at low energies

$$\frac{e^2}{4\pi} \simeq \frac{1}{137}$$

photons, leptons
nearly decouple

$$\frac{g^2}{4\pi} \simeq 1$$

gluons, quarks
confined

- Nuclear forces = van der Waals forces of QCD

3. Chiral symmetry

- Photons are extremely useful to probe QCD

Much of what we know about the structure of the hadrons stems from scattering experiments involving electrons or photons

$e + N \rightarrow e + N$ form factors of the nucleon

$e + N \rightarrow e + \textit{hadrons}$ deep inelastic scattering

electroproduction, photoproduction

- For bound states of quarks,
e.m. interaction is a small perturbation

Perturbation series in powers of $\frac{e^2}{4\pi}$ ✓

Discuss only the leading term: set $e = 0$

- Lagrangian then reduces to QCD

$$g, m_u, m_d, m_s, m_c, m_b, m_t$$

- m_u, m_d, m_s happen to be light

Consequence:

Approximate flavour symmetries

Play a crucial role for the low energy properties

Theoretical paradise

$$m_u = m_d = m_s = 0$$

$$m_c = m_b = m_t = \infty$$

QCD with 3 massless quarks

- Lagrangian contains a single parameter: g
 g is net colour of a quark
depends on radius of the region considered

- Colour contained within radius r

$$\frac{g^2}{4\pi} = \frac{2\pi}{9 |\ln(r \Lambda_{\text{QCD}})|}$$

- Intrinsic scale Λ_{QCD} is meaningful,
but not dimensionless

⇒ No dimensionless free parameter

All dimensionless physical quantities are pure numbers, determined by the theory

Cross sections can be expressed in terms of Λ_{QCD} or in the mass of the proton

- Interactions of u, d, s are identical
If the masses are set equal to zero,
there is no difference at all

$$q = \begin{pmatrix} u \\ d \\ s \end{pmatrix}$$

- Lagrangian symmetric under $u \leftrightarrow d \leftrightarrow s$

$$q' = V \cdot q \quad V \in \text{SU}(3)$$

V acts on quark flavour, mixes u, d, s

- More symmetry: For massless fermions,
right and left do not communicate
- ⇒ Lagrangian of massless QCD is invariant under independent rotations of the right- and left-handed quark fields

$$q_R = \frac{1}{2}(1 + \gamma_5) q, \quad q_L = \frac{1}{2}(1 - \gamma_5) q$$

$$q'_R = V_R \cdot q_R \quad q'_L = V_L \cdot q_L$$

$$\text{SU}(3)_R \times \text{SU}(3)_L$$

- Massless QCD invariant under $SU(3)_R \times SU(3)_L$

$SU(3)$ has 8 parameters

⇒ Symmetry under Lie group with 16 parameters

⇒ 16 conserved “charges”

Q_1^V, \dots, Q_8^V (vector currents, $R + L$)

Q_1^A, \dots, Q_8^A (axial currents, $R - L$)

commute with the Hamiltonian:

$$[Q_i^V, H_0] = 0 \quad [Q_i^A, H_0] = 0$$

“Chiral symmetry” of massless QCD

- Vafa and Witten 1984: state of lowest energy is invariant under the vector charges

$$Q_i^V |0\rangle = 0$$

- Axial charges ? $Q_i^A |0\rangle = ?$

Two alternatives for axial charges

$$Q_i^A |0\rangle = 0$$

Wigner-Weyl realization of G
ground state is symmetric

$$\langle 0 | \bar{q}_R q_L | 0 \rangle = 0$$

ordinary symmetry
spectrum contains parity partners
degenerate multiplets of G

$$Q_i^A |0\rangle \neq 0$$

Nambu-Goldstone realization of G
ground state is asymmetric

$$\langle 0 | \bar{q}_R q_L | 0 \rangle \neq 0$$

“order parameter”
spontaneously broken symmetry
spectrum contains Nambu-Goldstone bosons
degenerate multiplets of $SU(3)_V \subset G$

$$G = SU(3)_R \times SU(3)_L$$

- Spontaneous symmetry breakdown was discovered in condensed matter physics:

Spontaneous magnetization selects direction

⇒ Rotation symmetry is spontaneously broken

Nambu-Goldstone bosons = spin waves, magnons

- Nambu 1960: state of lowest energy in particle physics is not invariant under chiral rotations

$$Q_i^A |0\rangle \neq 0$$

For dynamical reasons, the state of lowest energy must be asymmetric

⇒ Chiral symmetry is spontaneously broken

- Very strong experimental evidence ✓
- Theoretical understanding on the basis of the QCD Lagrangian ?

- Analog of Magnetization ?

$$\bar{q}_R q_L = \begin{pmatrix} \bar{u}_R u_L & \bar{d}_R u_L & \bar{s}_R u_L \\ \bar{u}_R d_L & \bar{d}_R d_L & \bar{s}_R d_L \\ \bar{u}_R s_L & \bar{d}_R s_L & \bar{s}_R s_L \end{pmatrix}$$

Transforms like $(\bar{3}, 3)$ under $SU(3)_R \times SU(3)_L$

If the ground state were symmetric, the matrix $\langle 0 | \bar{q}_R q_L | 0 \rangle$ would have to vanish, because it singles out a direction in flavour space

“quark condensate”, is quantitative measure of spontaneous symmetry breaking

“order parameter”

$$\langle 0 | \bar{q}_R q_L | 0 \rangle \Leftrightarrow \text{magnetization}$$

- Ground state is invariant under $SU(3)_V$

$\Rightarrow \langle 0 | \bar{q}_R q_L | 0 \rangle$ is proportional to unit matrix

$$\langle 0 | \bar{u}_R u_L | 0 \rangle = \langle 0 | \bar{d}_R d_L | 0 \rangle = \langle 0 | \bar{s}_R s_L | 0 \rangle$$

$$\langle 0 | \bar{u}_R d_L | 0 \rangle = \dots = 0$$

4. Goldstone Theorem

- Consequence of $Q_i^A |0\rangle \neq 0$:

$$H_0 Q_i^A |0\rangle = Q_i^A H_0 |0\rangle = 0$$

spectrum must contain 8 states

$$Q_1^A |0\rangle, \dots, Q_8^A |0\rangle \quad \text{with } E = 0,$$

spin 0, negative parity, octet of $SU(3)_V$

Nambu-Goldstone bosons

- Argument is not water tight:

$$\langle 0 | Q_i^A Q_k^A | 0 \rangle = \int d^3x d^3y \langle 0 | A_i^0(x) A_k^0(y) | 0 \rangle$$

$$\langle 0 | A_i^0(x) A_k^0(y) | 0 \rangle \text{ only depends on } \vec{x} - \vec{y}$$

$\Rightarrow \langle 0 | Q_i^A Q_k^A | 0 \rangle$ is proportional to the volume of the universe, $|Q_i^A |0\rangle| = \infty$

- Rigorous version of Goldstone theorem:

$$\langle 0 | \bar{q}_R q_L | 0 \rangle \neq 0 \Rightarrow \exists \text{ massless particles}$$

Proof

fasten seatbelts: takes 3 slides

$$Q = \int d^3x \bar{u} \gamma^0 \gamma_5 d$$

$$[Q, \bar{d} \gamma_5 u] = -\bar{u}u - \bar{d}d$$

- $F^\mu(x - y) \equiv \langle 0 | \bar{u}(x) \gamma^\mu \gamma_5 d(x) \bar{d}(y) \gamma_5 u(y) | 0 \rangle$

Lorentz invariance $\Rightarrow F^\mu(z) = z^\mu f(z^2)$

Chiral symmetry $\Rightarrow \partial_\mu F^\mu(z) = 0$

$$F^\mu(z) = \frac{z^\mu}{z^4} \times \text{constant (for } z^2 \neq 0)$$

- Spectral decomposition:

$$\langle 0 | \bar{u}(x) \gamma^\mu \gamma_5 d(x) \bar{d}(y) \gamma_5 u(y) | 0 \rangle =$$

$$\sum_n \langle 0 | \bar{u} \gamma^\mu \gamma_5 d | n \rangle \langle n | \bar{d} \gamma_5 u | 0 \rangle e^{-i p_n(x-y)}$$

$p_n^0 \geq 0 \Rightarrow F^\mu(z)$ is analytic in z^0 for $\text{Im } z^0 < 0$

$$F^\mu(z) = \frac{z^\mu}{\{(z^0 - i\epsilon)^2 - \vec{z}^2\}^2} \times \text{constant}$$

- Positive frequency part of massless propagator: (exercise # 1)

$$\Delta^+(z, 0) = \frac{i}{(2\pi)^3} \int \frac{d^3p}{2p^0} e^{-ipz} \quad , \quad p^0 = |\vec{p}|$$

$$= \frac{1}{4\pi i \{(z^0 - i\epsilon)^2 - \vec{z}^2\}}$$

- Result

$$\langle 0 | \bar{u}(x) \gamma^\mu \gamma_5 d(x) \bar{d}(y) \gamma_5 u(y) | 0 \rangle = C \partial^\mu \Delta^+(z, 0)$$

- Compare Källén–Lehmann representation:

$$\langle 0 | \bar{u}(x) \gamma^\mu \gamma_5 d(x) \bar{d}(y) \gamma_5 u(y) | 0 \rangle$$

$$= (2\pi)^{-3} \int d^4p p^\mu \rho(p^2) e^{-ip(x-y)}$$

$$= \int_0^\infty ds \rho(s) \partial^\mu \Delta^+(x-y, s)$$

$\Delta^+(z, s) \iff$ massive propagator

$$\Delta^+(z, s) = \frac{i}{(2\pi)^3} \int d^4p \theta(p^0) \delta(p^2 - s) e^{-ipz}$$

\Rightarrow Only massless intermediate states contribute:

$$\rho(s) = C \delta(s)$$

- Why only massless intermediate states ?

$\langle n | \bar{d} \gamma_5 u | 0 \rangle \neq 0$ only if $\langle n |$ has spin 0

If $|n\rangle$ has spin 0 $\Rightarrow \langle 0 | \bar{u}(x) \gamma^\mu \gamma_5 d(x) | n \rangle \propto p^\mu e^{-ipx}$

$\partial_\mu (\bar{u} \gamma^\mu \gamma_5 d) = 0 \Rightarrow p^2 = 0$

\Rightarrow Either \exists massless particles or $C = 0$

- Claim: $\langle 0 | \bar{q}_R q_L | 0 \rangle \neq 0 \Rightarrow C \neq 0$

Lorentz invariance, chiral symmetry

$\Rightarrow \langle 0 | \bar{d}(y) \gamma_5 u(y) \bar{u}(x) \gamma^\mu \gamma_5 d(x) | 0 \rangle = C' \partial^\mu \Delta^-(z)$

$\Rightarrow \langle 0 | [\bar{u}(x) \gamma^\mu \gamma_5 d(x), \bar{d}(y) \gamma_5 u(y)] | 0 \rangle$

$$= C \partial^\mu \Delta^+(z, 0) - C' \partial^\mu \Delta^-(z, 0)$$

- Causality: if $x - y$ is spacelike, then

$\langle 0 | [\bar{u}(x) \gamma^\mu \gamma_5 d(x), \bar{d}(y) \gamma_5 u(y)] | 0 \rangle = 0$

$\Rightarrow C' = -C$

$\Rightarrow \langle 0 | [\bar{u}(x) \gamma^\mu \gamma_5 d(x), \bar{d}(y) \gamma_5 u(y)] | 0 \rangle = C \partial^\mu \Delta(z, 0)$

$\Rightarrow \langle 0 | [Q, \bar{d}(y) \gamma_5 u(y)] | 0 \rangle = C$

- $\langle 0 | [Q, \bar{d}(y) \gamma_5 u(y)] | 0 \rangle = -\langle 0 | \bar{u}u + \bar{d}d | 0 \rangle = C$

Hence $\langle 0 | \bar{u}u + \bar{d}d | 0 \rangle \neq 0$ implies $C \neq 0$ qed.

5. Gell-Mann-Oakes-Renner relation

- ⇒ Spectrum of QCD with 3 massless quarks must contain 8 massless physical particles, $J^P = 0^-$
- Indeed, the 8 lightest mesons do have these quantum numbers:

$$\pi^+, \pi^0, \pi^-, K^+, K^0, \bar{K}^0, K^-, \eta$$

But massless they are not

???

- Real world \neq paradise

$$m_u, m_d, m_s \neq 0$$

Quark masses break chiral symmetry,
allow the left to talk to the right

- Chiral symmetry broken in two ways:

spontaneously

$$\langle 0 | \bar{q}_R q_L | 0 \rangle \neq 0$$

explicitly

$$m_u, m_d, m_s \neq 0$$

- H_{QCD} only has approximate symmetry, to the extent that m_u, m_d, m_s are small

$$H_{\text{QCD}} = H_0 + H_1$$

$$H_1 = \int d^3x \{m_u \bar{u}u + m_d \bar{d}d + m_s \bar{s}s\}$$

- H_0 is Hamiltonian of the massless theory, invariant under $SU(3)_R \times SU(3)_L$
- H_1 breaks the symmetry, transforms with $(3, \bar{3}) \oplus (\bar{3}, 3)$
- For the low energy structure of QCD, the heavy quarks do not play an essential role: c, b, t are singlets under $SU(3)_R \times SU(3)_L$
Can include the heavy quarks in H_0
- Nambu-Goldstone bosons are massless only if the symmetry is exact

- Gell-Mann-Oakes-Renner relation:

$$M_\pi^2 = (m_u + m_d) \times |\langle 0 | \bar{u} u | 0 \rangle| \times \frac{1}{F_\pi^2}$$

1968

↑

explicit

↑

spontaneous

Coefficient: decay constant F_π

Derivation

takes 2 slides

- Pion matrix elements in massless theory:

$$\langle 0 | \bar{u} \gamma^\mu \gamma_5 d | \pi^- \rangle = i \sqrt{2} F p^\mu$$

$$\langle 0 | \bar{u} i \gamma_5 d | \pi^- \rangle = \sqrt{2} G$$

Only the one-pion intermediate state

$$\langle 0 | \bar{u}(x) \gamma^\mu \gamma_5 d(x) \overset{\uparrow}{\bar{d}(y) \gamma_5 u(y)} | 0 \rangle = C \partial^\mu \Delta^+(z, 0)$$

$$|\pi^- \rangle \langle \pi^- |$$

contributes. Hence $2 F G = C$

- Value of C fixed by quark condensate

$$C = -\langle 0 | \bar{u} u + \bar{d} d | 0 \rangle$$

⇒ Exact result in massless theory:

$$F G = -\langle 0 | \bar{u} u | 0 \rangle$$

- Matrix elements for $m_{\text{quark}} \neq 0$:

$$\langle 0 | \bar{u} \gamma^\mu \gamma_5 d | \pi^- \rangle = i \sqrt{2} F_\pi p^\mu$$

$$\langle 0 | \bar{u} i \gamma_5 d | \pi^- \rangle = \sqrt{2} G_\pi$$

- Current conservation

$$\partial_\mu (\bar{u} \gamma^\mu \gamma_5 d) = (m_u + m_d) \bar{u} i \gamma_5 d$$

$$\Rightarrow F_\pi M_\pi^2 = (m_u + m_d) G_\pi$$

$$\boxed{M_\pi^2 = (m_u + m_d) \frac{G_\pi}{F_\pi}} \quad \text{exact for } m \neq 0$$

- $F_\pi \rightarrow F$, $G_\pi \rightarrow G$ for $m \rightarrow 0$

$$F G = -\langle 0 | \bar{u} u | 0 \rangle$$

$$\Rightarrow \frac{G_\pi}{F_\pi} = -\frac{\langle 0 | \bar{u} u | 0 \rangle}{F_\pi^2} + O(m)$$

$$\Rightarrow M_\pi^2 = (m_u + m_d) \left(\frac{-\langle 0 | \bar{u} u | 0 \rangle}{F_\pi^2} \right) + O(m^2) \quad \checkmark$$

$$\Rightarrow \langle 0 | \bar{u} u | 0 \rangle \leq 0 \text{ if quark masses are positive}$$

- $M_\pi^2 = (m_u + m_d) B + O(m^2)$

$$B = \frac{|\langle 0 | \bar{u} u | 0 \rangle|}{F_\pi^2} \Big|_{m_u, m_d \rightarrow 0}$$

- M_π disappears if the symmetry breaking is turned off, $m_u, m_d \rightarrow 0$ ✓

- Explains why the pseudoscalar mesons have very different masses

$$M_{K^+}^2 = (m_u + m_s) B + O(m^2)$$

$$M_{K^0}^2 = (m_d + m_s) B + O(m^2)$$

⇒ M_K^2 is about 13 times larger than M_π^2 , because m_u, m_d happen to be small compared to m_s

- First order perturbation theory also yields


$$M_\eta^2 = \frac{1}{3} (m_u + m_d + 4m_s) B + O(m^2)$$

⇒ $M_\pi^2 - 4M_K^2 + 3M_\eta^2 = O(m^2)$

Gell-Mann-Okubo formula for M^2 ✓

Checking the GMOR formula on a lattice

- Can determine M_π as function of $m_u = m_d = m$



Lüscher, Lattice conference 2005 ETM collaboration, hep-lat/0701012

- No quenching, quark masses sufficiently light
- ⇒ Legitimate to use χ PPT for the extrapolation to the physical values of m_u, m_d

- Quality of data is impressive
- Proportionality of M_π^2 to the quark mass appears to hold out to values of m_u, m_d that are an order of magnitude larger than in nature
- Main limitation: systematic uncertainties
in particular: $N_f = 2 \rightarrow N_f = 3$

II. Chiral perturbation theory

Scholarpedia: *Chiral Perturbation Theory*

6. Group geometry

- QCD with 3 massless quarks:
spontaneous symmetry breakdown
from $SU(3)_R \times SU(3)_L$ to $SU(3)_V$
generates 8 Nambu-Goldstone bosons
- Generalization: suppose symmetry group
of Hamiltonian is Lie group G
Generators $Q_1, Q_2, \dots, Q_D, D = \dim(G)$
For some generators $Q_i |0\rangle \neq 0$
How many Nambu-Goldstone bosons ?
- Consider those elements of the Lie algebra
 $Q = \alpha_1 Q_1 + \dots + \alpha_n Q_D$, for which $Q |0\rangle = 0$
These elements form a subalgebra:
 $Q |0\rangle = 0, Q' |0\rangle = 0 \Rightarrow [Q, Q'] |0\rangle = 0$
Dimension of subalgebra: $d \leq D$
- Of the D vectors $Q_i |0\rangle$
 $D - d$ are linearly independent
 $\Rightarrow D - d$ different physical states of zero mass
 $\Rightarrow D - d$ Nambu-Goldstone bosons

- Subalgebra generates subgroup $H \subset G$
 H is symmetry group of the ground state
 coset space G/H contains as many parameters
 as there are Nambu-Goldstone bosons
 $d = \dim(H)$, $D = \dim(G)$

⇒ Nambu-Goldstone bosons live on the coset G/H

- Example: QCD with N_f massless quarks

$$G = \text{SU}(N_f)_R \times \text{SU}(N_f)_L$$

$$H = \text{SU}(N_f)_V$$

$$D = 2(N_f^2 - 1), \quad d = N_f^2 - 1$$

$$N_f^2 - 1 \text{ Nambu-Goldstone bosons}$$

- It so happens that $m_u, m_d \ll m_s$
- $m_u = m_d = 0$ is an excellent approximation
 $\text{SU}(2)_R \times \text{SU}(2)_L$ is a nearly exact symmetry
 $N_f = 2$, $N_f^2 - 1 = 3$ Nambu-Goldstone bosons
 (pions)

7. Generating functional of QCD

- Basic objects for quantitative analysis of QCD: Green functions of the currents

$$V_a^\mu = \bar{q} \gamma^\mu \frac{1}{2} \lambda_a q, \quad A_a^\mu = \bar{q} \gamma^\mu \gamma_5 \frac{1}{2} \lambda_a q,$$

$$S_a = \bar{q} \frac{1}{2} \lambda_a q, \quad P_a = \bar{q} i \gamma_5 \frac{1}{2} \lambda_a q$$

Include singlets, with $\lambda_0 = \sqrt{2/3} \times 1$, as well as

$$\omega = \frac{1}{16\pi^2} \text{tr}_c G_{\mu\nu} \tilde{G}^{\mu\nu}$$

- Can collect all of the Green functions formed with these operators in a generating functional: Perturb the system with external fields

$$v_\mu^a(x), a_\mu^a(x), s_a(x), p^a(x), \theta(x)$$

Replace the Lagrangian of the massless theory

$$\mathcal{L}_0 = -\frac{1}{2g^2} \text{tr}_c G_{\mu\nu} G^{\mu\nu} + \bar{q} i \gamma^\mu (\partial_\mu - i G_\mu) q$$

by $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_1$

$$\mathcal{L}_1 = v_\mu^a V_a^\mu + a_\mu^a A_a^\mu - s^a S_a - p^a P_a - \theta \omega$$

- Quark mass terms are included in the external field $s_a(x)$

- $|0 \text{ in}\rangle$: system is in ground state for $x^0 \rightarrow -\infty$
Probability amplitude for finding ground state when $x^0 \rightarrow +\infty$:

$$e^{iS_{\text{QCD}}\{v,a,s,p,\theta\}} = \langle 0 \text{ out} | 0 \text{ in}\rangle_{v,a,s,p,\theta}$$

- Expressed in terms of ground state of \mathcal{L}_0 :

$$e^{iS_{\text{QCD}}\{v,a,s,p,\theta\}} = \langle 0 | T \exp i \int dx \mathcal{L}_1 | 0 \rangle$$

- Expansion of $S_{\text{QCD}}\{v, a, s, p, \theta\}$ in powers of the external fields yields the connected parts of the Green functions of the massless theory

$$S_{\text{QCD}}\{v, a, s, p, \theta\} = - \int dx s_a(x) \langle 0 | S^a(x) | 0 \rangle \\ + \frac{i}{2} \int dx dy a_\mu^a(x) a_\nu^b(y) \langle 0 | T A_a^\mu(x) A_b^\nu(y) | 0 \rangle_{\text{conn}} + \dots$$

- $S_{\text{QCD}}\{v, a, s, p, \theta\}$ is referred to as the *generating functional* of QCD

- For Green functions of full QCD, set

$$s_a(x) = m_a + \tilde{s}_a(x), \quad m_a = \text{tr} \lambda_a m$$

and expand around $\tilde{s}_a(x) = 0$

- Path integral representation for generating functional:

$$e^{iS_{\text{QCD}}\{v,a,s,p\}} = \mathcal{N} \int [dG] e^{i \int dx \mathcal{L}_G} \det D$$

$$\mathcal{L}_G = -\frac{1}{2g^2} \text{tr}_c G_{\mu\nu} G^{\mu\nu} - \frac{\theta}{16\pi^2} \text{tr}_c G_{\mu\nu} \tilde{G}^{\mu\nu}$$

$$D = i\gamma^\mu \{ \partial_\mu - i(G_\mu + v_\mu + a_\mu \gamma_5) \} - s - i\gamma_5 p$$

G_μ is matrix in colour space

v_μ, a_μ, s, p are matrices in flavour space

$v_\mu(x) \equiv \frac{1}{2} \lambda_a v_\mu^a(x)$, etc.

8. Ward identities

Symmetry in terms of Green functions

- Lagrangian is invariant under

$$q_R(x) \rightarrow V_R(x) q_R(x), \quad q_L(x) \rightarrow V_L(x) q_L(x)$$
$$V_R(x), V_L(x) \in U(3)$$

provided the external fields are transformed with

$$v'_\mu + a'_\mu = V_R(v_\mu + a_\mu)V_R^\dagger - i\partial_\mu V_R V_R^\dagger$$
$$v'_\mu - a'_\mu = V_L(v_\mu - a_\mu)V_L^\dagger - i\partial_\mu V_L V_L^\dagger$$
$$s' + ip' = V_R(s + ip)V_L^\dagger$$

The operation takes the Dirac operator into

$$D' = \{P_- V_R + P_+ V_L\} D \{P_+ V_R^\dagger + P_- V_L^\dagger\}$$
$$P_\pm = \frac{1}{2}(1 \pm \gamma_5)$$

- $\det D$ requires regularization

∄ symmetric regularization

$$\Rightarrow \det D' \neq \det D, \text{ only } |\det D'| = |\det D|$$

symmetry does not survive quantization

- Change in $\det D$ can explicitly be calculated

For an infinitesimal transformation

$$V_R = 1 + i\alpha + i\beta + \dots, \quad V_L = 1 + i\alpha - i\beta + \dots$$

the change in the determinant is given by

$$\det D' = \det D e^{-i \int dx \{2\langle\beta\rangle\omega + \langle\beta\Omega\rangle\}}$$

$$\langle A \rangle \equiv \text{tr } A$$

$$\omega = \frac{1}{16\pi^2} \text{tr}_c G_{\mu\nu} \tilde{G}^{\mu\nu} \quad \text{gluons}$$

$$\Omega = \frac{N_c}{4\pi^2} \epsilon^{\mu\nu\rho\sigma} \partial_\mu v_\nu \partial_\rho v_\sigma + \dots \quad \text{ext. fields}$$

- Consequence for generating functional:

The term with ω amounts to a change in θ ,
can be compensated by $\theta' = \theta - 2\langle\beta\rangle$

Pull term with $\langle\beta\Omega\rangle$ outside the path integral

$$\Rightarrow S_{\text{QCD}}\{v', a', s', p', \theta'\} = S_{\text{QCD}}\{v, a, s, p, \theta\} - \int dx \langle\beta\Omega\rangle$$

$$S_{\text{QCD}}\{v', a', s', p', \theta'\} = S_{\text{QCD}}\{v, a, s, p, \theta\} - \int dx \langle \beta \Omega \rangle$$

- S_{QCD} is invariant under $U(3)_R \times U(3)_L$, except for a specific change due to the anomalies
- Relation plays key role in low energy analysis: collects all of the Ward identities
For the octet part of the axial current, e.g.

$$\begin{aligned} \partial_\mu^x \langle 0 | T A_a^\mu(x) P_b(y) | 0 \rangle &= -\frac{1}{4} i \delta(x - y) \langle 0 | \bar{q} \{ \lambda_a, \lambda_b \} q | 0 \rangle \\ &+ \langle 0 | T \bar{q}(x) i \gamma_5 \{ m, \frac{1}{2} \lambda_a \} q(x) P_b(y) | 0 \rangle \end{aligned}$$

- Symmetry of the generating functional implies the operator relations

$$\partial_\mu V_a^\mu = \bar{q} i [m, \frac{1}{2} \lambda_a] q, \quad a = 0, \dots, 8$$

$$\partial_\mu A_a^\mu = \bar{q} i \gamma_5 \{ m, \frac{1}{2} \lambda_a \} q, \quad a = 1, \dots, 8$$

$$\partial_\mu A_0^\mu = \sqrt{\frac{2}{3}} \bar{q} i \gamma_5 m q + \sqrt{6} \omega$$

- Textbook derivation of the Ward identities goes in inverse direction, but is slippery formal manipulations, anomalies ?

9. Low energy expansion

- If the spectrum has an energy gap
- ⇒ no singularities in scattering amplitudes or Green functions near $p = 0$
- ⇒ low energy behaviour may be analyzed with Taylor series expansion in powers of p


$$f(t) = 1 + \frac{1}{6} \langle r^2 \rangle t + \dots \text{ form factor}$$
$$T(p) = a + b p^2 + \dots \text{ scattering amplitude}$$

Cross section dominated by S -wave scattering length $\frac{d\sigma}{d\Omega} \simeq |a|^2$

- Expansion parameter: $\frac{p}{m} = \frac{\text{momentum}}{\text{energy gap}}$
- Taylor series only works if the spectrum has an energy gap, i.e. if there are no massless particles

- Illustration: Coulomb scattering

$$e + e \rightarrow e + e$$



Photon exchange \Rightarrow pole at $t = 0$

$$T = \frac{e^2}{(p' - p)^2}$$

Scattering amplitude does not admit Taylor series expansion in powers of p

- QCD does have an energy gap but the gap is very small: M_π
- \Rightarrow Taylor series has very small radius of convergence, useful only for $p < M_\pi$

- Massless QCD contains infrared singularities due to the Nambu-Goldstone bosons
 - For $m_u = m_d = 0$, pion exchange gives rise to poles and branch points at $p = 0$
- ⇒ Low energy expansion is not a Taylor series, contains logarithms

Singularities due to Nambu-Goldstone bosons can be worked out with an effective field theory

Chiral Perturbation Theory

Weinberg, Dashen, Pagels, Gasser, . . .

- Chiral perturbation theory correctly reproduces the infrared singularities of QCD
- Quantities of interest are expanded in powers of external momenta and quark masses
- Expansion has been worked out to next-to-leading order for many quantities
"Chiral perturbation theory to one loop"
- In quite a few cases, the next-to-next-to-leading order is also known

- Properties of the Nambu-Goldstone bosons are governed by the hidden symmetry that is responsible for their occurrence
- Focus on the singularities due to the pions

$$H_{\text{QCD}} = H_0 + H_1$$


$$H_1 = \int d^3x \{m_u \bar{u}u + m_d \bar{d}d\}$$

H_0 is invariant under $G = \text{SU}(2)_R \times \text{SU}(2)_L$

$|0\rangle$ is invariant under $H = \text{SU}(2)_V$

mass term of strange quark is included in H_0

- Treat H_1 as a perturbation



- Extension to $\text{SU}(3)_R \times \text{SU}(3)_L$ straightforward: include singularities due to exchange of K, η
- \Rightarrow Discuss this later, first treat only m_u, m_d as small quantities, keep m_s fixed at the physical value, study the effective theory belonging to $\text{SU}(2)_R \times \text{SU}(2)_L$

10. Effective Lagrangian

- Replace quarks and gluons by pions

$$\vec{\pi}(x) = \{\pi^1(x), \pi^2(x), \pi^3(x)\}$$

$$\mathcal{L}_{\text{QCD}} \rightarrow \mathcal{L}_{\text{eff}}$$

- Central claim:

A. Effective theory yields alternative representation for generating functional of QCD

$$e^{iS_{\text{QCD}}\{v,a,s,p,\theta\}} = \mathcal{N}_{\text{eff}} \int [d\pi] e^{i \int dx \mathcal{L}_{\text{eff}}\{\vec{\pi},v,a,s,p,\theta\}}$$

B. \mathcal{L}_{eff} has the same symmetries as \mathcal{L}_{QCD}

⇒ Can calculate the low energy expansion of the Green functions with the effective theory.

If \mathcal{L}_{eff} is chosen properly, this reproduces the low energy expansion of QCD, order by order.

- Proof of A and B: H.L., Annals Phys. 1994

- Pions live on the coset $G/H = \text{SU}(2)$

$$\vec{\pi}(x) \rightarrow U(x) \in \text{SU}(2)$$

The fields $\vec{\pi}(x)$ are the coordinates of $U(x)$

Can use canonical coordinates, for instance

$$U = \exp i \vec{\pi} \cdot \vec{\tau} \in \text{SU}(2)$$

- Action of the symmetry group on the quarks:

$$q'_R = V_R \cdot q_R, \quad q'_L = V_L \cdot q_L$$

- Action on the pion field:

$$U' = V_R \cdot U \cdot V_L^\dagger$$

Note: Transformation law for the coordinates $\vec{\pi}$ is complicated, nonlinear

- Except for the contribution from the anomalies, \mathcal{L}_{eff} is invariant

$$\mathcal{L}_{eff}\{U', v', a', s', p', \theta'\} = \mathcal{L}_{eff}\{U, v, a, s, p, \theta\}$$

Symmetry of S_{QCD} implies symmetry of \mathcal{L}_{eff}

Side remark

- For nonrelativistic effective theories, the effective Lagrangian is in general invariant only up to a total derivative.
- ⇒ From the point of view of effective field theory, nonrelativistic systems with Nambu-Goldstone bosons are more complicated than relativistic ones

detailed discussion: H. L., Phys. Rev. D49 (1994) 3033

- Origin of the complication: the generators of the symmetry group may themselves give rise to order parameters

$$\langle 0 | Q^i | 0 \rangle \neq 0$$

This cannot happen in the relativistic case:

$$Q = \int d^3x j^0(x)$$
$$\langle 0 | j^\mu(x) | 0 \rangle = 0 \Rightarrow \langle 0 | Q | 0 \rangle = 0$$

Nonrelativistic example where it does happen:

Heisenberg model of a ferromagnet

$$H = -g \sum_{\langle ij \rangle} \vec{s}_i \cdot \vec{s}_j$$

$g > 0$ $\uparrow\uparrow$ lower in energy than $\uparrow\downarrow$

- Ground state = $\uparrow\uparrow\uparrow\uparrow \cdots \uparrow\uparrow$
- Magnetization: $\vec{M} = \frac{\mu}{V} \sum_i \vec{s}_i$
 $\langle 0 | \vec{M} | 0 \rangle \neq 0 \iff \langle 0 | \bar{q}_R q_L | 0 \rangle \neq 0$
- Symmetry generators: $\vec{Q} = \sum_i \vec{s}_i \propto \vec{M}$
- Hamiltonian is invariant under the full rotation group $G = \text{SO}(3)$, ground state is invariant only under rotations around the direction of $\langle 0 | \vec{M} | 0 \rangle$, $H = \text{U}(1)$
- Effective field lives on $G/H = S_2$: unit vector \vec{U} , parametrized by 2 coordinates π^1, π^2 .
- Effective Lagrangian of ferromagnet is invariant under local rotations only up to a total derivative. Leading term is related to the Brouwer degree of the map $(\pi^1, \pi^2) \rightarrow \vec{U}$.

11. Explicit construction of \mathcal{L}_{eff}

- First ignore the external fields,

$$\mathcal{L}_{eff} = \mathcal{L}_{eff}(U, \partial U, \partial^2 U, \dots)$$

Derivative expansion:

$$\mathcal{L}_{eff} = f_0(U) + f_1(U) \times \square U + f_2(U) \times \partial_\mu U \times \partial^\mu U + \dots$$

$\uparrow \qquad \qquad \uparrow \qquad \qquad \uparrow$
 $O(1) \qquad O(p^2) \qquad O(p^2)$

Amounts to expansion in powers of momenta

- Term of $O(1)$: $f_0(U) = f_0(V_R U V_L^\dagger)$

$$V_R = \mathbf{1}, \quad V_L = U \rightarrow V_R U V_L^\dagger = \mathbf{1}$$

$\Rightarrow f_0(U) = f_0(\mathbf{1})$ irrelevant constant, drop it

- Term with $\square U$: integrate by parts

\Rightarrow can absorb $f_1(U)$ in $f_2(U)$

⇒ Derivative expansion of \mathcal{L}_{eff} starts with

$$\mathcal{L}_{eff} = f_2(U) \times \partial_\mu U \times \partial^\mu U + O(p^4)$$

- Replace the partial derivative by

$$\Delta_\mu \equiv \partial_\mu U U^\dagger, \quad \text{tr} \Delta_\mu = 0$$

Δ_μ is invariant under $SU(2)_L$ and transforms with the representation $D^{(1)}$ under $SU(2)_R$:

$$\Delta_\mu \rightarrow V_R \Delta_\mu V_R^\dagger$$

In this notation, leading term is of the form

$$\mathcal{L}_{eff} = \tilde{f}_2(U) \times \Delta_\mu \times \Delta^\mu + O(p^4)$$

- Invariance under $SU(2)_L$: $\tilde{f}_2(U) = \tilde{f}_2(U V_L^\dagger)$
- ⇒ $\tilde{f}_2(U)$ is independent of U
- Invariance under $SU(2)_R$: $\Delta_\mu \times \Delta^\mu$ transforms with $D^{(1)} \times D^{(1)} \rightarrow$ contains unity exactly once: $\text{tr}(\Delta_\mu \Delta^\mu) = \text{tr}(\partial_\mu U U^\dagger \partial^\mu U U^\dagger) = -\text{tr}(\partial_\mu U \partial^\mu U^\dagger)$
- ⇒ Geometry fixes leading term up to a constant

$$\mathcal{L}_{eff} = \frac{F^2}{4} \text{tr}(\partial_\mu U \partial^\mu U^\dagger) + O(p^4)$$

$$\mathcal{L}_{eff} = \frac{F^2}{4} \text{tr}(\partial_\mu U \partial^\mu U^\dagger) + O(p^4)$$

- Lagrangian of the nonlinear σ -model
- Expansion in powers of $\vec{\pi}$:

$$U = \exp i \vec{\pi} \cdot \vec{\tau} = 1 + i \vec{\pi} \cdot \vec{\tau} - \frac{1}{2} \vec{\pi}^2 + \dots$$

$$\Rightarrow \mathcal{L}_{eff} = \frac{F^2}{2} \partial_\mu \vec{\pi} \cdot \partial^\mu \vec{\pi} + \frac{F^2}{48} \text{tr}\{[\partial_\mu \pi, \pi] [\partial^\mu \pi, \pi]\} + \dots$$

For the kinetic term to have the standard normalization: rescale the pion field, $\vec{\pi} \rightarrow \vec{\pi}/F$

$$\mathcal{L}_{eff} = \frac{1}{2} \partial_\mu \vec{\pi} \cdot \partial^\mu \vec{\pi} + \frac{1}{48F^2} \text{tr}\{[\partial_\mu \pi, \pi] [\partial^\mu \pi, \pi]\} + \dots$$

- \Rightarrow a. Symmetry requires the pions to interact
- b. Derivative coupling: Nambu-Goldstone bosons only interact if their momentum does not vanish $\Rightarrow \cancel{\lambda} \pi^4$

- Expression given for \mathcal{L}_{eff} only holds if the external fields are turned off. Also, $\text{tr}(\partial_\mu U \partial^\mu U^\dagger)$ is invariant only under global transformations
- Suffices to replace $\partial_\mu U$ by

$$D_\mu U = \partial_\mu U - i(v_\mu + a_\mu)U + iU(v_\mu - a_\mu)$$

In contrast to $\text{tr}(\partial_\mu U \partial^\mu U^\dagger)$, the term $\text{tr}(D_\mu U D^\mu U^\dagger)$ is invariant under local $SU(2)_R \times SU(2)_L$

- Can construct further invariants: $s + ip$ transforms like $U \Rightarrow \text{tr}\{(s + ip)U^\dagger\}$ is invariant
Violates parity, but $\text{tr}\{(s + ip)U^\dagger\} + \text{tr}\{(s - ip)U\}$ is even under $p \rightarrow -p, \vec{\pi} \rightarrow -\vec{\pi}$
- In addition, \exists invariant independent of U :
 $D_\mu \theta D^\mu \theta$, with $D_\mu \theta = \partial_\mu \theta + 2 \text{tr}(a_\mu)$
- Count the external fields as
 $\theta = O(1), \quad v_\mu, a_\mu = O(p), \quad s, p = O(p^2)$

- Derivative expansion yields string of the form

$$\mathcal{L}_{eff} = \mathcal{L}^{(2)} + \mathcal{L}^{(4)} + \mathcal{L}^{(6)} + \dots$$

- Full expression for leading term:

$$\mathcal{L}^{(2)} = \frac{F^2}{4} \langle D_\mu U D^\mu U^\dagger + \chi U^\dagger + U \chi^\dagger \rangle + h_0 D_\mu \theta D^\mu \theta$$

$$\chi \equiv 2B(s + ip), \quad \langle X \rangle \equiv \text{tr}(X)$$

- At LO, symmetry allows 2 "low energy constants" (F, B) plus 1 "contact term" (h_0)
- Next-to-leading order:

$$\begin{aligned} \mathcal{L}^{(4)} = & \frac{\ell_1}{4} \langle D_\mu U D^\mu U^\dagger \rangle^2 + \frac{\ell_2}{4} \langle D_\mu U D_\nu U^\dagger \rangle \langle D^\mu U D^\nu U^\dagger \rangle \\ & + \frac{\ell_3}{4} \langle \chi U^\dagger + U \chi^\dagger \rangle^2 + \frac{\ell_4}{4} \langle D_\mu \chi D^\mu U^\dagger + D_\mu U D^\mu \chi^\dagger \rangle \\ & + \dots \end{aligned}$$

- Altogether 7 LEC + 3 CT at NLO
- Number of LEC rapidly grows with the order of the expansion

- Infinitely many LEC
Symmetry does not determine these
Predictivity ?
- Essential point: If \mathcal{L}_{eff} is known to given order
 \Rightarrow can work out low energy expansion of the
Green functions to that order Weinberg 1979
- F_π, M_π involve 2 LEC at NLO: l_3, l_4 .
- In the $\pi\pi$ scattering amplitude, two further
LEC enter at NLO: l_1, l_2 .
- Note: effective theory is a quantum field theory
Need to perform the path integral

$$e^{iS_{\text{QCD}}\{v,a,s,p,\theta\}} = \mathcal{N}_{eff} \int [dU] e^{i \int dx \mathcal{L}_{eff}\{U,v,a,s,p,\theta\}}$$

- Classical theory \Leftrightarrow tree graphs
Need to include graphs with loops
- Power counting in dimensional regularization:
Graphs with ℓ loops are suppressed by factor $p^{2\ell}$ as compared to tree graphs
- \Rightarrow Leading contributions given by tree graphs
Graphs with one loop contribute at next-to-leading order, etc.
- The leading contribution to S_{QCD} is given by the sum of all tree graphs = classical action:

$$S_{\text{QCD}}\{v, a, s, p, \theta\} = \underset{U(x)}{\text{extremum}} \int dx \mathcal{L}_{\text{eff}}\{U, v, a, s, p, \theta\}$$

III. Illustrations

12. Some tree level calculations

12.1 Extracting the quark condensate from the generating functional

$$e^{iS_{\text{QCD}}\{v,a,s,p,\theta\}} = \langle 0 | T \exp i \int dx \mathcal{L}_1 | 0 \rangle$$

- To calculate the quark condensate of the massless theory, it suffices to consider the generating functional for $v = a = p = \theta = 0$ and to take a constant scalar external field

$$s = \begin{pmatrix} m_u & 0 \\ 0 & m_d \end{pmatrix}$$

$$\mathcal{L}_1 = -(m_u \bar{u}u + m_d \bar{d}d)$$

$$S_{\text{QCD}}^1 = - \int dx \langle 0 | m_u \bar{u}u + m_d \bar{d}d | 0 \rangle$$

- $\Rightarrow \langle 0 | \bar{u}u | 0 \rangle$ and $\langle 0 | \bar{d}d | 0 \rangle$ are the coefficients of the terms in S_{QCD} that are linear in m_u and m_d

12.2 Condensate in terms of effective theory

- Need the effective action for $v = a = p = \theta = 0$ to first order in

$$s = \begin{pmatrix} m_u & 0 \\ 0 & m_d \end{pmatrix}$$

⇒ classical level of effective theory suffices.

- extremum of the classical action: $U = 1$

$$S_{\text{QCD}}^1 = \int dx \langle F^2 B s \rangle = \int dx F^2 B (m_u + m_d)$$

- comparison with

$$S_{\text{QCD}}^1 = - \int dx \langle 0 | m_u \bar{u}u + m_d \bar{d}d | 0 \rangle \quad \text{yields}$$

$$\boxed{\langle 0 | \bar{u}u | 0 \rangle = \langle 0 | \bar{d}d | 0 \rangle = -F^2 B} \quad (1)$$

Quark condensate in chiral limit:


$$\Sigma = |\langle 0 | \bar{u}u | 0 \rangle|_{m_u, m_d \rightarrow 0}$$

$$\Sigma = F^2 B$$

Lattice result (FLAG 2013, arXiv:1310.8555):

$$N_f = 2 : \quad \Sigma = 270(7) \text{ MeV}$$

$$N_f = 2 + 1 : \quad \Sigma = 265(17) \text{ MeV}$$



12.3 Evaluation of M_π at tree level

- In classical theory, the square of the mass is the coefficient of the term in the Lagrangian that is quadratic in the meson field:

$$\begin{aligned}\frac{F^2}{4}\langle\chi U^\dagger + U\chi^\dagger\rangle &= \frac{F^2 B}{2}\langle m(U^\dagger + U)\rangle \\ &= F^2 B(m_u + m_d)\left\{1 - \frac{\vec{\pi}^2}{2F^2} + \dots\right\}\end{aligned}$$

Hence
$$M_\pi^2 = (m_u + m_d)B \quad (2)$$

- Tree level result for F_π :

$$F_\pi = F \quad (3)$$

- (1) + (2) + (3) \Rightarrow GMOR relation:

$$M_\pi^2 = \frac{(m_u + m_d) |\langle 0 | \bar{u}u | 0 \rangle|}{F_\pi^2}$$

13. M_π beyond tree level


- The formula $M_\pi^2 = (m_u + m_d)B$ only holds at tree level, represents leading term in expansion of M_π^2 in powers of m_u, m_d
- Disregard isospin breaking: set $m_u = m_d = m$

13.1 M_π to 1 loop

- Claim: at next-to-leading order, the expansion of M_π^2 in powers of m contains a logarithm:

$$M_\pi^2 = M^2 - \frac{1}{2} \frac{M^4}{(4\pi F)^2} \ln \frac{\Lambda_3^2}{M^2} + O(M^6)$$
$$M^2 \equiv 2mB$$

- Proof: Pion mass \Leftrightarrow pole position, for instance in the Fourier transform of $\langle 0 | T A_a^\mu(x) A_b^\nu(y) | 0 \rangle$
Suffices to work out the perturbation series for this object to one loop of the effective theory



- Result (exercise # 5):

$$M_\pi^2 = M^2 + \frac{2\ell_3 M^4}{F^2} + \frac{M^2}{2F^2} \frac{1}{i} \Delta(0, M^2) + O(M^6)$$

$\Delta(0, M^2)$ is the propagator at the origin
(exercise # 2):

$$\begin{aligned} \Delta(0, M^2) &= \frac{1}{(2\pi)^d} \int \frac{d^d p}{M^2 - p^2 - i\epsilon} \\ &= i (4\pi)^{-d/2} \Gamma(1 - d/2) M^{d-2} \end{aligned}$$

- Contains a pole at $d = 4$:

$$\Gamma(1 - d/2) = \frac{2}{d - 4} + \dots$$

- Divergent part is proportional to M^2 :

$$\begin{aligned} M^{d-2} &= M^2 \mu^{d-4} (M/\mu)^{d-4} = M^2 \mu^{d-4} e^{(d-4) \ln(M/\mu)} \\ &= M^2 \mu^{d-4} \{1 + (d-4) \ln(M/\mu) + \dots\} \end{aligned}$$

- Denote the singular factor by

$$\begin{aligned}\lambda &\equiv \frac{1}{2}(4\pi)^{-d/2} \Gamma(1 - d/2) \mu^{d-4} \\ &= \frac{\mu^{d-4}}{16\pi^2} \left\{ \frac{1}{d-4} - \frac{1}{2}(\ln 4\pi + \Gamma'(1) + 1) + O(d-4) \right\}\end{aligned}$$

- The propagator at the origin then becomes

$$\frac{1}{i}\Delta(0, M^2) = M^2 \left\{ 2\lambda + \frac{1}{16\pi^2} \ln \frac{M^2}{\mu^2} + O(d-4) \right\}$$

- In the expression for M_π^2

$$M_\pi^2 = M^2 + \frac{2\ell_3 M^4}{F^2} + \frac{M^2}{2F^2} \frac{1}{i}\Delta(0, M^2) + O(M^6)$$

the divergence can be absorbed in ℓ_3 :

$$\ell_3 = -\frac{1}{2}\lambda + \ell_3^{\text{ren}}$$

- ℓ_3^{ren} depends on the renormalization scale μ

$$\ell_3^{\text{ren}} = \frac{1}{64\pi^2} \ln \frac{\mu^2}{\Lambda_3^2} \quad \text{running low energy constant}$$

- Λ_3 is the ren. group invariant scale of ℓ_3

- Net result for M_π^2

$$M_\pi^2 = M^2 - \frac{1}{2} \frac{M^4}{(4\pi F)^2} \ln \frac{\Lambda_3^2}{M^2} + O(M^6)$$

⇒ M_π^2 contains a chiral logarithm at NLO

- Crude estimate for Λ_3 , based on SU(3) mass formulae for the pseudoscalar octet:

$$0.2 \text{ GeV} < \Lambda_3 < 2 \text{ GeV}$$

$$\bar{\ell}_3 \equiv \ln \frac{\Lambda_3^2}{M_\pi^2} = 2.9 \pm 2.4$$

Gasser, L. 1984

∃ better determination $\bar{\ell}_3$ on the lattice, to be discussed later

⇒ Next-to-leading term is small correction:

$$0.005 < \frac{1}{2} \frac{M_\pi^2}{(4\pi F_\pi)^2} \ln \frac{\Lambda_3^2}{M_\pi^2} < 0.040$$

- Scale of the expansion is set by size of pion mass in units of decay constant:

$$\frac{M^2}{(4\pi F)^2} \simeq \frac{M_\pi^2}{(4\pi F_\pi)^2} = 0.0144$$

13.2 M_π to 2 loops

- Terms of order m_{quark}^3 :

$$M_\pi^2 = M^2 - \frac{1}{2} \frac{M^4}{(4\pi F)^2} \ln \frac{\Lambda_3^2}{M^2} + \frac{17}{18} \frac{M^6}{(4\pi F)^4} \left(\ln \frac{\Lambda_M^2}{M^2} \right)^2 + k_M M^6 + O(M^8)$$

F is pion decay constant for $m_u = m_d = 0$

ChPT to two loops

Colangelo 1995

- Coefficients $\frac{1}{2}$ and $\frac{17}{18}$ determined by symmetry
- Λ_3, Λ_M and $k_M \iff$ LEC in \mathcal{L}_{eff}

14. F_π to one loop

- Also contains a logarithm at NLO:

$$F_\pi = F \left\{ 1 - \frac{M^2}{16\pi^2 F^2} \ln \frac{M^2}{\Lambda_4^2} + O(M^4) \right\}$$
$$M_\pi^2 = M^2 \left\{ 1 + \frac{M^2}{32\pi^2 F^2} \ln \frac{M^2}{\Lambda_3^2} + O(M^4) \right\}$$

F is pion decay constant in limit $m_u, m_d \rightarrow 0$

- Structure is the same, coefficients and scale of logarithm are different
- Low energy theorem: at leading order in the chiral expansion, the *scalar radius* is also determined by the scale Λ_4 :

$$\langle r^2 \rangle_s = \frac{6}{(4\pi F)^2} \left\{ \ln \frac{\Lambda_4^2}{M^2} - \frac{13}{12} + O(M^2) \right\}$$

Chiral symmetry relates F_π to $\langle r^2 \rangle_s$

What is the scalar radius ? \Rightarrow next section

15. Pion form factors

- Scalar form factor of the pion:

$$F_s(t) = \langle \pi(p') | \bar{q} q | \pi(p) \rangle, \quad t = (p' - p)^2$$

- Definition of scalar radius:

$$F_s(t) = F_s(0) \left\{ 1 + \frac{1}{6} \langle r^2 \rangle_s t + O(t^2) \right\}$$

- Low energy theorem:

$$\langle r^2 \rangle_s = \frac{6}{(4\pi F)^2} \left\{ \ln \frac{\Lambda_4^2}{M^2} - \frac{13}{12} + O(M^2) \right\}$$

⇒ In massless QCD, the scalar radius diverges, because the density of the pion cloud only decreases with a power of the distance

- Same infrared singularity also occurs in the charge radius (e.m. current), but coefficient of the chiral logarithm is 6 times smaller:

$$\langle r^2 \rangle_s = \frac{6}{(4\pi F)^2} \left\{ \ln \frac{\Lambda_4^2}{M^2} - \frac{13}{12} + O(M^2) \right\}$$

$$\langle r^2 \rangle_{em} = \frac{1}{(4\pi F)^2} \left\{ \ln \frac{\Lambda_6^2}{M^2} - 1 + O(M^2) \right\}$$

⇒ $\langle r^2 \rangle_s > \langle r^2 \rangle_{em}$ if M small enough

- $\langle r^2 \rangle_{em}$ can be determined experimentally

$$\langle r^2 \rangle_{em} = 0.439 \pm 0.008 \text{ fm}^2$$

NA7 Collaboration, NP B277 (1986) 168

- Scalar form factor of the pion can be calculated by means of dispersion theory
- Result for the slope:

$$\langle r^2 \rangle_s = 0.61 \pm 0.04 \text{ fm}^2$$

Colangelo, Gasser, L., Nucl. Phys. 2001

⇒ Corresponding value of the scale Λ_4 :

$$\Lambda_4 = 1.26 \pm 0.14 \text{ GeV}$$


16. Lattice results for M_π, F_π

16.1 Results for M_π

- Determine the scale Λ_3 by comparing the lattice results for M_π as function of m with the χ PT formula

$$M_\pi^2 = M^2 - \frac{1}{2} \frac{M^4}{(4\pi F)^2} \ln \frac{\Lambda_3^2}{M^2} + O(M^6)$$

$$M^2 \equiv 2Bm$$



lattice results for \bar{l}_3

Horizontal axis shows the value of $\bar{l}_3 \equiv \ln \frac{\Lambda_3^2}{M_\pi^2}$


Range for Λ_3 obtained in 1984 corresponds to $\bar{l}_3 = 2.9 \pm 2.4$

	$N_f = 2$	$N_f = 2 + 1$	$N_f = 2 + 1 + 1$
\bar{l}_3	3.45 ± 0.26	2.77 ± 1.27	3.70 ± 0.27

FLAG 2013

16.2 Results for F_π

$$F_\pi = F \left\{ 1 - \frac{M^2}{16\pi^2 F^2} \ln \frac{M^2}{\Lambda_4^2} + O(M^4) \right\}$$



Horizontal axis shows the value of $\bar{\ell}_4 \equiv \ln \frac{\Lambda_4^2}{M_\pi^2}$

- Lattice results beautifully confirm the prediction for the sensitivity of F_π to m_u, m_d :

$$\frac{F_\pi}{F} = 1.072 \pm 0.007$$

Colangelo, Dürr 2004

17. $\pi\pi$ scattering

17.1 Low energy scattering of pions

- Consider scattering of pions with $\vec{p} = 0$
 - At $\vec{p} = 0$, only the S-waves survive (angular momentum barrier). Moreover, these reduce to the scattering lengths
 - Bose statistics: S-waves cannot have $I = 1$, either have $I = 0$ or $I = 2$
- ⇒ At $\vec{p} = 0$, the $\pi\pi$ scattering amplitude is characterized by two constants: a_0^0, a_0^2
- Chiral symmetry suppresses the interaction at low energy: Nambu-Goldstone bosons of zero momentum do not interact
- ⇒ a_0^0, a_0^2 disappear in the limit $m_u, m_d \rightarrow 0$
- ⇒ $a_0^0, a_0^2 \sim M_\pi^2$ measure symmetry breaking

17.2 Tree level of χ PT

- Low Energy theorem Weinberg 1966:

$$a_0^0 = \frac{7M_\pi^2}{32\pi F_\pi^2} + O(M_\pi^4)$$
$$a_0^2 = -\frac{M_\pi^2}{16\pi F_\pi^2} + O(M_\pi^4)$$

\Rightarrow Chiral symmetry predicts a_0^0, a_0^2 in terms of F_π

- Accuracy is limited: Low energy theorem only specifies the first term in the expansion in powers of the quark masses
Corrections from higher orders ?

17.3 Scattering lengths at 1 loop

- Next term in the chiral perturbation series:

$$a_0^0 = \frac{7M_\pi^2}{32\pi F_\pi^2} \left\{ 1 + \frac{9}{2} \frac{M_\pi^2}{(4\pi F_\pi)^2} \ln \frac{\Lambda_0^2}{M_\pi^2} + O(M_\pi^4) \right\}$$

- Coefficient of chiral logarithm unusually large
Strong, attractive final state interaction
- Scale Λ_0 is determined by the LEC of $\mathcal{L}_{eff}^{(4)}$:

$$\frac{9}{2} \ln \frac{\Lambda_0^2}{M_\pi^2} = \frac{20}{21} \bar{\ell}_1 + \frac{40}{21} \bar{\ell}_2 - \frac{5}{14} \bar{\ell}_3 + 2\bar{\ell}_4 + \frac{5}{2}$$

- Information about $\bar{\ell}_1, \dots, \bar{\ell}_4$?


$\bar{\ell}_1, \bar{\ell}_2 \iff$	momentum dependence of scattering amplitude
-----------------------------------	--

\Rightarrow Can be determined phenomenologically

$\bar{\ell}_3, \bar{\ell}_4 \iff$	dependence of scattering amplitude on quark masses
-----------------------------------	---

Have discussed their values already


17.4 Numerical predictions from χ P T



Sizable corrections in a_0^0 | a_0^2 nearly stays put

17.5 a_0^0, a_0^2 from lattice results for l_3, l_4

- Uncertainty in prediction for a_0^0, a_0^2 is dominated by the uncertainty in the LEC l_3, l_4
- Can make use of the lattice results for these



17.6 Experiments concerning a_0^0, a_0^2

- Production experiments $\pi N \rightarrow \pi\pi N$,
 $\psi \rightarrow \pi\pi\omega$, $B \rightarrow D\pi\pi$, ...

Problem: pions are not produced in vacuo

⇒ Extraction of $\pi\pi$ scattering amplitude is not simple

Accuracy rather limited


- $K^\pm \rightarrow \pi^+\pi^-e^\pm\nu$ data:
CERN-Saclay, E865, NA48/2
- $K^\pm \rightarrow \pi^0\pi^0\pi^\pm$, $K^0 \rightarrow \pi^0\pi^0\pi^0$: cusp near threshold, NA48/2
- $\pi^+\pi^-$ atoms, DIRAC

17.7 Results from K_{e4} decay

$$K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu$$

- Allows clean measurement of $\delta_0^0 - \delta_1^1$

Theory predicts $\delta_0^0 - \delta_1^1$ as function of energy




Prediction: $a_0^0 = 0.220 \pm 0.005$

NA48/2: $a_0^0 = 0.2206 \pm 0.0049 \pm 0.0018 \pm 0.0064$
stat *syst* *theo*

Bloch-Devaux, Chiral Dynamics 2009


- There was a discrepancy here, because a pronounced isospin breaking effect from $K \rightarrow \pi^0 \pi^0 e \nu \rightarrow \pi^+ \pi^- e \nu$ had not been accounted for in the data analysis

Colangelo, Gasser, Rusetsky 2007, Bloch-Devaux 2007



- The correction is not enormous, but matters: If a_0^0 is determined from the uncorrected NA48 data, the central value comes out higher than the theoretical prediction by about 4 times the uncertainty attached to this prediction.

17.8 Summary for a_0^0, a_0^2



18. Conclusions for $SU(2) \times SU(2)$

- Expansion in powers of m_u, m_d yields a very accurate low energy representation of QCD
- Lattice results confirm the GMOR relation
- ⇒ M_π is dominated by the contribution from the quark condensate
- ⇒ Energy gap of QCD is understood very well
- Lattice approach allows an accurate measurement of the low energy constant l_3 already now
- Even for l_4 , the lattice starts becoming competitive with dispersion theory

19. Expansion in powers of m_s

- The χ PT formulae for the expansion of many quantities of physical interest in powers of m_u , m_d , m_s have been worked out to NNLO, not only masses and decay constants, also form factors, $\eta \rightarrow 3\pi$,
- Theoretical reasoning:
 - Pion physics: expansion in powers of m_u, m_d works very well.
 - Physics of the strange particles: $SU(3)_V$ is an approximate symmetry.
 - \Rightarrow Symmetry breaking parameter $m_s - m_{ud}$ must be small, meaningful to expand in powers of $m_s - m_{ud}$.
 - Since $m_u, m_d \ll m_s$
 - $\Rightarrow m_s$ can be treated as a perturbation
 - \Rightarrow Expect expansion in powers of m_s to work, but convergence to be comparatively slow
- I do not know of an alternative explanation of the empirical fact that $SU(3)$ is an approximate symmetry.

19.1 Form of the effective Lagrangian

- If all three light quark masses vanish, the QCD Lagrangian is invariant under $SU(3)_R \times SU(3)_L$
- The spontaneous breakdown to $SU(3)_V$ generates eight Nambu-Goldstone bosons:

$$\pi^\pm, \pi^0, K^\pm, K^0, \bar{K}^0, \eta$$

⇒ Effective fields can be collected in $U \in SU(3)$

$$U(x) = \exp i \pi(x)$$

$$\pi(x) = \lambda_1 \pi^1(x) + \dots + \lambda_8 \pi^8(x)$$

- Symmetry again fixes the leading term in \mathcal{L}_{eff} :

$$\mathcal{L}^{(2)} = \frac{F_0^2}{4} \langle D_\mu U D^\mu U^\dagger + \chi U^\dagger + U \chi^\dagger \rangle + \frac{H_0}{12} D_\mu \theta D^\mu \theta$$

$$\chi \equiv 2 B_0 (s + ip), \quad \langle X \rangle \equiv \text{tr}(X)$$

- U is now 3x3, χ is 3x3, otherwise the form of the Lagrangian is the same as for $SU(2)_R \times SU(2)_L$:
- Symmetry does not determine F_0, B_0, H_0

$$\mathcal{L}^{(2)} = \frac{F_0^2}{4} \langle D_\mu U D^\mu U^\dagger + \chi U^\dagger + U \chi^\dagger \rangle + \frac{H_0}{12} D_\mu \theta D^\mu \theta$$

$$\chi \equiv 2 B_0 (s + ip), \quad \langle X \rangle \equiv \text{tr}(X)$$

- Significance of F_0, B_0 : leading terms in the expansion of the decay constants and meson masses in powers of the quark masses:

$$\langle 0 | \bar{u} \gamma_\mu \gamma_5 d | \pi^- \rangle = \sqrt{2} F_0 p^\mu \{1 + O(m)\}$$

$$\langle 0 | \bar{u} \gamma_\mu \gamma_5 s | K^- \rangle = \sqrt{2} F_0 p^\mu \{1 + O(m)\}$$

$$\langle 0 | \bar{d} \gamma_\mu \gamma_5 s | K^0 \rangle = \sqrt{2} F_0 p^\mu \{1 + O(m)\}$$

$$M_{\pi^-}^2 = (m_u + m_d) B_0 + O(m^2)$$

$$M_{K^-}^2 = (m_u + m_s) B_0 + O(m^2)$$

$$M_{K^0}^2 = (m_d + m_s) B_0 + O(m^2)$$

- The expansion includes m_s
 - $\Rightarrow F_0, B_0$ are independent of m_u, m_d, m_s
 - In the effective theory built on $SU(2)_R \times SU(2)_L$ m_s is not an expansion parameter
 - $\Rightarrow F, B$ do depend on m_s
- $$F_0 = F|_{m_s \rightarrow 0} \quad B_0 = B|_{m_s \rightarrow 0}$$

- Next-to-leading order:

$$\begin{aligned}
\mathcal{L}^{(4)} = & L_1 \langle D_\mu U D^\mu U^\dagger \rangle^2 + L_2 \langle D_\mu U D_\nu U^\dagger \rangle \langle D^\mu U D^\nu U^\dagger \rangle \\
& + L_3 \langle D_\mu U D^\mu U^\dagger D_\nu U D^\nu U^\dagger \rangle + L_4 \langle D_\mu U D^\mu U^\dagger \rangle \langle \chi U^\dagger + U \chi^\dagger \rangle \\
& + L_5 \langle D_\mu U D^\mu U^\dagger (\chi U^\dagger + U \chi^\dagger) \rangle + L_6 \langle \chi U^\dagger + U \chi^\dagger \rangle^2 \\
& + L_7 \langle \chi U^\dagger - U \chi^\dagger \rangle^2 + L_8 \langle \chi U^\dagger \chi U^\dagger + U \chi^\dagger U \chi^\dagger \rangle \\
& - iL_9 \langle F_{\mu\nu}^R D^\mu U D^\nu U^\dagger + F_{\mu\nu}^L D^\mu U^\dagger D^\nu U \rangle + L_{10} \langle F_{\mu\nu}^R U F^{\mu\nu L} U^\dagger \rangle \\
& + H_1 \langle F_{\mu\nu}^R F^{\mu\nu R} + F_{\mu\nu}^L F^{\mu\nu L} \rangle + H_2 \langle \chi \chi^\dagger \rangle
\end{aligned}$$

10 LEC + 2 CT: $L_1, \dots, L_{10}; H_1, H_2$

$\Rightarrow m_u, m_d, m_s, F_0, B_0, L_4, L_5, L_6, L_7, L_8$ determine

$M_{\pi^\pm}, M_{\pi^0}, M_{K^\pm}, M_{K^0}, M_{\bar{K}^0}, M_\eta, F_{\pi^\pm}, F_{\pi^0}, F_{K^\pm}, F_{K^0}, F_{\bar{K}^0}, F_\eta$ to NLO


- compare SU(2)×SU(2):

7 LEC + 3 CT: $\ell_1, \dots, \ell_7; h_1, h_2, h_3$

$\Rightarrow m_u, m_d, F, B, \ell_3, \ell_4, \ell_7$ determine

$M_{\pi^\pm}, M_{\pi^0}, F_{\pi^\pm}, F_{\pi^0}$ to NLO

19.2 Results for L_4, L_5, L_6, L_8



Numerical values shown refer to running scale $\mu = M_\rho$

⇒ For PACS-CS, only the statistical errors are indicated

- The crude estimates given in 1985 for the LEC relevant at NLO are confirmed
- However: not all of the lattice data on the quark mass dependence of M_π, M_K, F_π, F_K are well described by the χ PT formulae

- m_s is not very small, terms of order m_s^2 yield sizable corrections.
 - Often, m_s is taken in the vicinity of the physical value while m_{ud} is significantly larger than the physical value
- ⇒ M_K, M_η are larger than the physical values, may be beyond reach
- The constants relevant at NNLO are still poorly known. Often, theoretical estimates are used, obtained by saturating sum rules with resonance contributions. Those constants that govern the dependence on the quark masses, however, represent integrals over scalar spectral functions. Scalar meson dominance does not work!
- ⇒ Theoretical estimates can at best indicate the order of magnitude.
- The lattice approach is the ideal method for the determination of the LEC !
 - Please do not use 'theory' for the LEC.

20. Zweig rule

- Concerns the role played by the sea quarks in physical matrix elements.

Okubo 1963, Zweig 1964, Iizuka 1966

- Leading low energy constants in the effective Lagrangian of $SU(2) \times SU(2)$: F and B

$$\{F, B, \Sigma\} = \left\{ F_\pi, \frac{M_\pi^2}{m_u + m_d}, |\langle 0 | \bar{u}u | 0 \rangle| \right\}_{m_u, m_d \rightarrow 0}$$

- Low energy theorem: $\Sigma = F^2 B$
exact, holds for any value of m_s .
- Zweig rule: F and B are independent of m_s .
- F_0, B_0, Σ_0 : values for $m_s = 0$
- *Paramagnetic inequalities*: both F and Σ decrease if m_s is taken smaller

$$F > F_0, \Sigma > \Sigma_0 \quad \text{Descotes-Genon, Girlanda \& Stern 2000}$$

- Behaviour if N_c becomes large:

F, B, Σ become independent of m_s if $N_c \rightarrow \infty$

$$F/F_0 \rightarrow 1, B/B_0 \rightarrow 1, \Sigma/\Sigma_0 \rightarrow 1$$

⇒ The differences $F/F_0 - 1, B/B_0 - 1, \Sigma/\Sigma_0 - 1$ measure the violations of the Zweig rule

- Expansion to NLO involves the low energy constants L_4 and L_6 of the $SU(3) \times SU(3)$ Lagrangian:


$$F/F_0 = 1 + \frac{8\bar{M}_K^2}{F_0^2} L_4 + \chi \log + \dots$$

$$\Sigma/\Sigma_0 = 1 + \frac{32\bar{M}_K^2}{F_0^2} L_6 + \chi \log + \dots$$

$$B/B_0 = 1 + \frac{16\bar{M}_K^2}{F_0^2} (2L_6 - L_4) + \chi \log + \dots$$

$$\bar{M}_K^2 \equiv m_s B_0$$

⇒ The LEC L_4 and L_6 measure the deviations from the Zweig rule



- The LEC appear to obey the Zweig-rule reasonably well: the values for L_4 , L_6 , $2L_6 - L_4$ are consistent with zero
- Lattice data leave much to be desired: only two papers without red tags in FLAG review: MILC (2009), HPQCD (2013).

- Inserting the lattice results for L_4, L_6 in the NLO formulae of χ PT, I get

	F/F_0	B/B_0	Σ/Σ_0
MILC (2009)	1.12(4)	1.10(7)	1.34(13)
HPQCD (2013)	1.10(8)	1.12(8)	1.32(28)
GL (1985)	1.0(1)	1.0(2)	1.0(3)

- ⇒ Evidence for small Zweig rule violations, consistent with the crude old estimates.

The Zweig rule violations roughly amount to a common change in scale:

$$F \simeq ZF_0 \quad B \simeq ZB_0 \quad \Rightarrow \quad \Sigma \simeq Z^3\Sigma_0$$

with $Z \simeq 1.10(5)$

- Paramagnetic inequalities of Descotes-Genon, Girlanda & Stern are confirmed.

- MILC has evaluated the ratios to all orders in m_s :

	F/F_0	B/B_0	Σ/Σ_0
NLO	1.12(4)	1.10(7)	1.34(13)
all orders	1.10(4)	1.20(7)	1.48(16)

For F/F_0 , the corrections are small, but for B/B_0 , the central values of the terms of order m_s and m_s^2 (or higher) are of the same size . . .

- ⇒ The Zweig rule deserves more attention !
- HPQCD instead evaluated the quark condensates at the physical quark masses:

$$\frac{\langle 0 | \bar{s}s | 0 \rangle}{\langle 0 | \bar{u}u | 0 \rangle} = 1.08(16)(1)$$

Confirms that SU(3) is a decent approximate symmetry: the symmetry breaking generated by $m_s - m_{ud}$ is too small to stick out from the noise of the calculation.

21. Quark mass ratios

21.1 Isospin breaking

- The symmetry properties of the vacuum shield the pions from isospin breaking:

The difference between m_u and m_d only generates a tiny effect of order

$$M_{\pi^+}^2 - M_{\pi^0}^2 \propto (m_u - m_d)^2.$$

- ⇒ The mass difference between π^0 and π^+ is due almost exclusively to electromagnetism.
- ⇒ More easy to determine the mean mass $m_{ud} \equiv \frac{1}{2}(m_u + m_d)$ than the difference $m_u - m_d$.
- Estimate the e.m. self-energies with the Dashen theorem:

$$M_{K^+}^2|_{e.m.} = M_{\pi^+}^2|_{e.m.} \quad M_{\pi^0}^2|_{e.m.} = M_{K^0}^2|_{e.m.} = 0$$

21.2 Quark mass ratios at leading order

- Solve the tree level mass formulae for the ratios m_s/m_{ud} and m_u/m_d : Weinberg 1977

$$\frac{m_s}{m_{ud}} = \frac{M_{K^+}^2 + M_{K^0}^2 - M_{\pi^+}^2}{M_{\pi^0}^2} = 25.9$$


$$\frac{m_u}{m_d} = \frac{M_{K^+}^2 - M_{K^0}^2 + 2M_{\pi^0}^2 - M_{\pi^+}^2}{M_{K^0}^2 - M_{K^+}^2 + M_{\pi^+}^2} = 0.56$$

- Low energy theorems, valid to leading order of the chiral expansion.

Corrections from higher orders ? Could they strongly modify the numerical result ?

What is the uncertainty to be attached to these predictions ?

21.3 Lattice results for m_s/m_{ud}



lattice average quoted in FLAG 2013:

$$\frac{m_s}{m_{ud}} = 27.46(15)(41)$$

$$27.46 = 25.9 + 1.6$$


\uparrow \uparrow

leading order higher orders

\Rightarrow correction is small, leading term of chiral perturbation series dominates

accuracy reached: 1.6 %

21.4 m_s/m_d versus m_u/m_d



- Most lattice calculations are done in pure QCD.
- For m_s/m_{ud} , this is a good approximation, because the uncertainties in the violations of the Dashen theorem do not strongly affect this ratio.
- For m_u/m_d , the situation is different. Lattice simulations of QCD + QED cannot be done with the same level of confidence as for QCD alone: not all systematic errors are under control (quenched photons, finite size effects for interactions of long range).

21.5 Low energy theorem valid to NLO

- The lattice result for m_s/m_{ud} determines the size of the correction in the relation

$$\frac{M_K^2}{M_\pi^2} = \frac{m_s + m_{ud}}{m_u + m_d} \left\{ 1 + \Delta_M \right\}$$

$$m_s/m_{ud} = 27.5 \pm 0.4 \Rightarrow \Delta_M = -0.057 \pm 0.013.$$

- Remarkably, chiral symmetry implies that the correction of NLO in the ratio of mass splittings is the same:

$$\frac{M_{K^0}^2 - M_{K^+}^2}{M_K^2 - M_\pi^2} = \frac{m_d - m_u}{m_s - m_{ud}} \left\{ 1 + \Delta_M + O(\mathcal{M}^2) \right\}$$

Hence the quark mass ratio

$$Q^2 \equiv \frac{m_s^2 - m_{ud}^2}{m_d^2 - m_u^2}$$

is given by a ratio of meson masses, up to corrections of NNLO:

$$Q^2 = \frac{M_K^2 - M_\pi^2}{M_{K^0}^2 - M_{K^+}^2} \cdot \frac{M_K^2}{M_\pi^2} \left\{ 1 + O(\mathcal{M}^2) \right\}$$

Gasser & L. 1985

21.6 Consequences for Q


- Insert Weinberg's leading order ratios

⇒ $Q = 24.3$.

- Q^2 is a ratio of quark mass squares

⇒ a given value of Q imposes a homogeneous quadratic constraint on m_u, m_d, m_s

⇒ represents an ellipse in the plane of the quark mass ratios:



- Critical input here is the "Dashen theorem": Weinberg's estimates for the quark mass ratios account for QED only to LO.

22. The decay $\eta \rightarrow 3\pi$

- The decay $\eta \rightarrow 3\pi$ provides a better handle on Q than the mass splitting between K^+ and K^0 , because the e.m. interaction is suppressed (Sutherland's theorem).
- For $e = 0$ and $m_u = m_d$, isospin is conserved, hence G-parity is conserved.

In this limit, the η is a stable particle:

$$G_\eta = 1, G_\pi = -1.$$

- ⇒ Since the e.m. contributions are tiny, the transition amplitude is to a very good approximation proportional to $(m_u - m_d)$.

22.1 Tree level

- Parameter free prediction for the leading term of the chiral perturbation series:

$$A(\eta \rightarrow \pi^+ \pi^- \pi^0) = -\frac{\sqrt{3}}{4} \cdot \frac{m_d - m_u}{m_s - m_{ud}} \cdot \frac{s - \frac{4}{3}M_\pi^2}{F_\pi^2}$$

- Compare leading term in the chiral expansion of the $\pi\pi$ scattering amplitude:

$$A(\pi\pi \rightarrow \pi\pi) = \frac{s - M_\pi^2}{F_\pi^2}$$

- In both cases, the leading term is linear in s and contains an Adler zero

$\pi\pi$ scattering

$$s_A = M_\pi^2$$

η decay

$$s_A = \frac{4}{3}M_\pi^2$$

- The analytic structure of the two amplitudes is very similar.
- In both cases, the higher order contributions of the chiral perturbation series are dominated by the final state interaction among the pions.

22.2 One loop

- Most remarkable property of the one loop representation: expressed in terms of F_π , F_K , M_π , M_K , M_η , Q , all LEC except L_3 drop out.

Gasser & L. 1985

$$A(\eta \rightarrow \pi^+ \pi^- \pi^0) = -\frac{1}{Q^2} \cdot \frac{M_K^2 (M_K^2 - M_\pi^2)}{3\sqrt{3} M_\pi^2 F_\pi^2} \cdot M(s, t, u)$$

- Moreover, L_3 concerns the momentum dependence of the amplitude, can be determined quite well from $\pi\pi$ scattering.
- ⇒ At one loop, the result for the rate is of the form

$$\Gamma_{\eta \rightarrow \pi^+ \pi^- \pi^0} = \frac{C}{Q^4} \quad Q^2 \equiv \frac{m_s^2 - m_{ud}^2}{m_d^2 - m_u^2}$$

where C is a known constant ⇒ Q can be determined from the observed rate.

- The main problem is not the uncertainty in L_3 , but the contributions from higher orders.

- In 1985, we estimated the uncertainty in the result for Q at

$$\frac{1}{Q^2} = (1.9 \pm 0.3) \cdot 10^{-3} \quad \Leftrightarrow \quad Q = 22.9_{-1.6}^{+2.1}$$

Gasser & L. 1985


- The result is consistent with the value $Q = 24.3$ obtained from the kaon mass difference with the Dashen theorem, but the uncertainties are large.

22.3 Dispersion theory

- The structure of the decay amplitude is governed by the final state interaction.

Standard method for the analysis of this interaction: dispersion theory.

- Main difference to $\pi\pi$ scattering: the subtraction constants relevant for $\eta \rightarrow 3\pi$ cannot be predicted to the same precision.
- Can analyze $\pi\pi$ scattering by treating only m_u and m_d as small: $SU(2) \times SU(2)$
- In η decay, need to treat m_s as an expansion parameter as well: $SU(3) \times SU(3)$
- Only the occurrence of an Adler zero follows from $SU(2) \times SU(2)$ symmetry alone.
- The subtraction constants can be estimated by comparing the dispersive and chiral representations at small values of s , t or u and requiring the occurrence of an Adler zero at the proper place.



Anisovich & L. 1996

⇒ Final state interaction amplifies the transition.

- Effect of the higher order contributions on the result for Q is modest:

$$Q = 22.4 \pm 0.9$$

Kambor, Wiesendanger & Wyler 1996

$$Q = 22.7 \pm 0.8$$

Anisovich & L. 1996

- Confirmed the one loop result, $Q = 22.9_{-1.6}^{+2.1}$, uncertainty reduced by a factor of 2.

22.4 Recent work on $\eta \rightarrow 3\pi$

- In the meantime, the experimental situation improved a lot: KLOE, MAMI, WASA.
- At low energies, the $\pi\pi$ phase shifts are now known to remarkable accuracy:
- Low energy precision experiments (E865, NA48, DIRAC).
- Low energy theorems for scattering lengths.
- Dispersion theory (Roy equations).
- χ PT has been worked out to NNLO.

Bijnens & Ghorbani 2007

- At the precision reached, isospin breaking needs to be accounted for.
- Nonrelativistic effective theory.

Ditsche, Kubis & Meissner 2009


Gullström, Kupsc & Rusetsky 2009

Schneider, Kubis & Ditsche 2011


- Improved dispersive analysis is under way

Colangelo, Lanz, L. & Passemar

- Preliminary results for the quark mass ratios



- Intersection moves to values of m_u/m_d and m_s/m_d that are somewhat smaller than those obtained with the LO mass formulae of Weinberg.



- The preliminary results for the ratio m_u/m_d are consistent with the lattice averages quoted by FLAG, but tend to be somewhat smaller.

23. Conclusions for $SU(3) \times SU(3)$

- Expansion in powers of m_s appears to work:

In all cases I know, where the calculation is under control, the truncation at low order yields a decent approximation

⇒ The picture looks coherent, also for $SU(3) \times SU(3)$

- $m_s \gg m_u, m_d \Rightarrow$ higher orders more important
- For many observables \exists representation to NNLO

Bijnens and collaborators

- Main problem: new LEC relevant at NNLO
- \exists estimates based on resonance models
- Vector meson dominance ✓
- Dependence on m_u, m_d, m_s : scalar resonances
- Scalar meson dominance ?
- Lattice results now start providing more precise values for the LEC, but the settling of dust is a slow process . . .

24. $T \neq 0$

- Most likely distribution of a given energy: thermal equilibrium, characterized by T
- Partition function: $Z = \text{Tr} e^{-\frac{H}{T}}$

24.1 Magnets

$$H = H_0 - \int d^3x \vec{H} \cdot \vec{M}$$

\vec{H} : external magnetic field


\vec{M} : magnetization

- Expectation value of magnetization:

$$\langle \vec{M} \rangle = \frac{1}{Z} \text{Tr} \left\{ e^{-\frac{H}{T}} \vec{M} \right\}$$

- $\langle \vec{M} \rangle$ is parallel to \vec{H}
- *Spontaneous magnetization:*

$\langle \vec{M} \rangle$ stays $\neq 0$ if $\vec{H} \rightarrow 0$



Temperature dependence of spontaneous magnetization


24.2 QCD

$$H = H_0 + \int d^3x \{m_u \bar{u}u + m_d \bar{d}d\}$$

- Partition function: $Z = \text{Tr} e^{-\frac{H}{T}}$
- Quark condensate at nonzero temperature:

$$\langle \bar{u}u \rangle = \frac{1}{Z} \text{Tr} \left\{ e^{-\frac{H}{T}} \bar{u}u \right\}$$

- For $T \rightarrow 0$, $\langle \bar{u}u \rangle$ tends to $\langle 0 | \bar{u}u | 0 \rangle$



- Symmetry relevant here: $SU(2)_R \times SU(2)_L$
Symmetry is exact only for $m_u, m_d \rightarrow 0$

24.3 Partition function of QCD at low T

- Insert complete set of states $H|n\rangle = E_n|n\rangle$

$$Z = \text{Tr} e^{-\frac{H}{T}} = \sum_n e^{-\frac{E_n}{T}}$$

- Only states with $E_n \lesssim T$ contribute

At $T = 0$ only the vacuum survives, $|n\rangle = |0\rangle$

- At low T, the next most important contribution stems from the pions
 - Pions of low energy behave like free particles, only interact weakly
- ⇒ At low energies, the partition function of QCD describes a gas of free pions

24.4 Partition function of free Bose gas

Bose 1924, Einstein 1925

- Complete set of one particle modes
- Label the modes with $k = 1, 2, 3, \dots$
- Example: box of size $L \times L \times L$, plane waves

$$\vec{p} = \frac{2\pi}{L} \{k_1, k_2, k_3\}, \quad k_r \in Z$$

$$k \leftrightarrow \{k_1, k_2, k_3\}$$

- bosons: $n_k = 0, 1, 2, \dots$ particles in each mode
- fermions: $n_k = 0$ or 1

⇒ Complete set of states for the entire gas:

$$|n\rangle = |n_1, n_2, \dots\rangle$$

- Energy in mode k : $\omega_k = \sqrt{m^2 + \vec{p}^2}$
- Energy of the gas in such a state

$$E_n = E_0 + n_1\omega_1 + n_2\omega_2 + \dots$$

↑ vacuum energy

$$\begin{aligned}
Z &= \sum_{n_1, n_2, \dots} e^{-\frac{1}{T}\{E_0 + n_1\omega_1 + n_2\omega_2 + \dots\}} \\
&= \sum_{n_1, n_2, \dots} e^{-\frac{E_0}{T}} \times e^{-\frac{n_1\omega_1}{T}} \times e^{-\frac{n_2\omega_2}{T}} \dots \\
&= e^{-\frac{E_0}{T}} \times \frac{1}{1 - e^{-\frac{\omega_1}{T}}} \times \frac{1}{1 - e^{-\frac{\omega_2}{T}}} \dots \\
&= e^{-\frac{E_0}{T}} \times \prod_k \frac{1}{1 - e^{-\frac{\omega_k}{T}}}
\end{aligned}$$

$$\ln Z = -\frac{E_0}{T} - \sum_k \ln(1 - e^{-\frac{\omega_k}{T}})$$

- Number of states in $\Delta^3 p$: $\frac{\Delta^3 p}{(2\pi/L)^3} = \frac{\Delta^3 p V}{(2\pi)^3}$

This reproduces a general rule of statistical mechanics: the volume element $\Delta^3 p \Delta^3 x$ of phase space contains $\Delta^3 p \Delta^3 x / h^3$ quantum states

$$\Rightarrow \ln Z = -\frac{E_0}{T} - \frac{V}{(2\pi)^3} \int d^3 p \ln(1 - e^{-\frac{\omega_{\vec{p}}}{T}})$$

24.5 Melting of the condensate

- QCD with two light flavours: 3 NGBs

$$\ln Z_{\text{QCD}} = -\frac{E_0}{T} - 3 \frac{V}{(2\pi)^3} \int d^3 p \ln(1 - e^{-\frac{\omega_{\vec{p}}}{T}}) + \dots$$

- Calculate the condensate from the partition function

$$\frac{\partial Z_{\text{QCD}}}{\partial m_u} = \frac{\partial \text{Tr} e^{-\frac{H}{T}}}{\partial m_u} = -\frac{1}{T} \text{Tr} \left\{ e^{-\frac{H}{T}} \frac{\partial H}{\partial m_u} \right\}$$

$$H = H_0 + \int d^3 x \{ m_u \bar{u} u + m_d \bar{d} d \}$$

$$\frac{\partial H}{\partial m_u} = \int d^3 x \bar{u} u$$

$$\begin{aligned} \frac{\partial Z_{\text{QCD}}}{\partial m_u} &= -\frac{1}{T} \int d^3 x \text{Tr} \left\{ e^{-\frac{H}{T}} \bar{u} u \right\} = -\frac{V}{T} \text{Tr} \left\{ e^{-\frac{H}{T}} \bar{u} u \right\} \\ &= -\frac{V}{T} \langle \bar{u} u \rangle Z_{\text{QCD}} \end{aligned}$$

$$\Rightarrow \langle \bar{u} u \rangle = -\frac{T}{V} \frac{\partial \ln Z_{\text{QCD}}}{\partial m_u}$$

$$\ln Z_{\text{QCD}} = -\frac{E_0}{T} - 3 \frac{V}{(2\pi)^3} \int d^3p \ln(1 - e^{-\frac{\omega_{\vec{p}}}{T}})$$

- First term dominates at low T:

$$\langle \bar{u}u \rangle = \frac{1}{V} \frac{\partial E_0}{\partial m_u} + \dots \text{ independent of } T$$

$$\frac{1}{V} \frac{\partial E_0}{\partial m_u} = \langle 0 | \bar{u}u | 0 \rangle$$

- Second term also depends on m_u , via

$$\omega_{\vec{p}} = \sqrt{M_\pi^2 + \vec{p}^2} \quad M_\pi^2 = (m_u + m_d)B + \dots$$

$$\Rightarrow \frac{\partial \omega_{\vec{p}}}{\partial m_u} = \frac{B}{2\omega_{\vec{p}}} = -\frac{\langle 0 | \bar{u}u | 0 \rangle}{2\omega_{\vec{p}} F^2}$$

$$\langle \bar{u}u \rangle = \langle 0 | \bar{u}u | 0 \rangle \left\{ 1 - \frac{3}{16\pi^3 F^2} \int \frac{d^3p}{\omega_{\vec{p}}} \frac{1}{\left(e^{\frac{\omega_{\vec{p}}}{T}} - 1 \right)} + \dots \right\}$$

$$\Rightarrow \langle \bar{u}u \rangle < \langle 0 | \bar{u}u | 0 \rangle \text{ quark condensate melts}$$

- Melting for massless quarks:


$$m_u, m_d \rightarrow 0 \Rightarrow M_\pi = 0 \Rightarrow \omega_{\vec{p}} = |\vec{p}|$$

$$\int \frac{d^3p}{|\vec{p}|} \frac{1}{\left(e^{\frac{|\vec{p}|}{T}} - 1\right)} = \frac{2\pi^3 T^2}{3}$$

$$\Rightarrow \langle \bar{u}u \rangle = \langle 0 | \bar{u}u | 0 \rangle \left\{ 1 - \frac{T^2}{8F^2} + O(T^4) \right\}$$

First two terms in temperature expansion of the quark condensate for $m_u = m_d = 0$. The constant F is the value of F_π in this limit.

Binétruy & Gaillard 1985



24.6 Pressure, energy density

- Free energy: $Z_{\text{QCD}} = e^{-\frac{F_{\text{QCD}}}{T}}$

$$F_{\text{QCD}} = E_0 + \frac{3VT}{(2\pi)^3} \int d^3p \ln(1 - e^{-\frac{\omega_{\vec{p}}}{T}}) + \dots$$

Free energy of noninteracting Bose gas
3 flavours: π^-, π^0, π^+

- Massless pions: $\int d^3p \ln(1 - e^{-\frac{|\vec{p}|}{T}}) = -\frac{4\pi^5}{45} T^3$

$$\Rightarrow F_{\text{QCD}} = E_0 - \frac{\pi^2}{30} VT^4 + O(T^6)$$

- $P_{\text{QCD}} = -\frac{\partial F_{\text{QCD}}}{\partial V} = \frac{\pi^2}{30} T^4 + O(T^6)$ pressure

- $s_{\text{QCD}} = \frac{\partial P_{\text{QCD}}}{\partial T} = \frac{2\pi^2}{15} T^3 + O(T^5)$ entropy density

- $u_{\text{QCD}} = Ts - P = \frac{\pi^2}{10} T^4 + O(T^6)$ energy density

\Rightarrow Leading terms in massless QCD are the same as for black body radiation, except for a factor 3/2 (3 independent pion states of a given momentum, 2 independent photon states)

24.7 Comparison: high temperature

- Asymptotic freedom \Rightarrow at high temperature, QCD also represents a gas of free particles: quarks, gluons
- High temperatures are beyond reach of χ PT
Instead: perturbation theory in powers of α_s
- Gas of free gluons: energy density, pressure, differ from the expressions obtained for a gas of free pions only by the factor $\frac{2 \times 8}{3}$
- Quarks are fermions, obey different statistics

- Partition function of free fermions

$$Z = \sum_{n_1, n_2, \dots} e^{-\frac{E_0}{T}} \times e^{-\frac{n_1 \omega_1}{T}} \times e^{-\frac{n_2 \omega_2}{T}} \dots$$

$$\begin{aligned} Z &= e^{-\frac{E_0}{T}} \times (1 + e^{-\frac{\omega_1}{T}}) \times (1 + e^{-\frac{\omega_2}{T}}) \dots \\ &= e^{-\frac{E_0}{T}} \times \prod_k (1 + e^{-\frac{\omega_k}{T}}) \end{aligned}$$

$$\ln Z = -\frac{E_0}{T} + \sum_k \ln(1 + e^{-\frac{\omega_k}{T}})$$

$$\Rightarrow \ln Z = -\frac{E_0}{T} + \frac{V}{(2\pi)^3} \int d^3 p \ln(1 + e^{-\frac{\omega_{\vec{p}}}{T}})$$

$$\Rightarrow \boxed{F = E_0 - \frac{VT}{(2\pi)^3} \int d^3 p \ln(1 + e^{-\frac{\omega_{\vec{p}}}{T}})}$$

Free energy of noninteracting fermions

$$\omega_{\vec{p}} = \sqrt{m^2 + \vec{p}^2}$$

- Massless quarks: $\omega_{\vec{p}} = |\vec{p}|$

$$\int d^3 p \ln(1 + e^{-\frac{|\vec{p}|}{T}}) = \frac{7\pi^5}{90} T^3$$

$$\Rightarrow F_{\text{quarks}} = E_0 - 3 \cdot 2 \cdot 2 \cdot N_f \frac{7\pi^2}{720} T^4 V$$

- Net result for energy density of QCD with N_f massless quarks

- High temperature:

$$u_{\text{QCD}} = \frac{\pi^2}{30} T^4 \left\{ \underset{\substack{\uparrow \\ \text{gluons}}}{8 \cdot 2} + \frac{7}{8} \cdot \underset{\substack{\uparrow \\ \text{quarks}}}{3 \cdot 2 \cdot 2} \cdot N_f \right\} + \dots$$

- Low temperature:

$$u_{\text{QCD}} = \frac{\pi^2}{30} T^4 \cdot \underset{\substack{\uparrow \\ \text{Nambu-Goldstone Bosons}}}{(N_f^2 - 1)} + \dots$$

- Contributions from other particles at low T:

$$\sim e^{-M_\rho/T} \simeq 0.006 \text{ for } \simeq 150 \text{ MeV}$$

- Probability to find a ρ is small, but:

many statistically independent states

⇒ Already at $T = 130 \text{ MeV}$, more energy is stored in $K, \eta, \rho \dots$ than in the pions

- Return to the quark condensate at low T
QCD with N_f massless quark flavours

$$\langle \bar{u}u \rangle = \langle 0 | \bar{u}u | 0 \rangle \left\{ 1 - \frac{(N_f^2 - 1) T^2}{12 N_f F^2} + O(T^4) \right\}$$

- Term of order T^4 ? Very tedious to do the calculation by hand, use the washing machine:

24.8 Chiral Perturbation Theory for $T \neq 0$

- Standard procedure:
 - Path integral representation for the transition amplitude in quantum mechanics

$$\langle x' | e^{-itH} | x \rangle = \int [Dx] e^{i \int_0^t dt' L}$$

Integration extends over all paths $x(t)$ with $x(0) = x$, $x(t) = x'$

- Path integral representation for the matrix elements of $e^{-\beta H}$: continue analytically in t to $t = -i\beta$

$$\langle x' | e^{-\beta H} | x \rangle = \int [Dx] e^{-\int_0^\beta dt' L^{eucl}}$$

- Partition function in this notation:

$$\text{Tr } e^{-\beta H} = \int dx \langle x | e^{-\beta H} | x \rangle \text{ with } \boxed{\beta = \frac{1}{T}}$$

⇒ Set $x' = x$ and integrate over x

$$\boxed{\text{Tr } e^{-\beta H} = \int [Dx] e^{-\int_0^\beta dt' L^{eucl}}}$$

Integration over all paths with $x(\beta) = x(0)$

- Formula also holds for path integral representation of QCD:

$$\text{Tr } e^{-\beta H} = \mathcal{N} \int [dG] e^{-\int_0^\beta dx^4 \int d^3x \mathcal{L}_G^{eucl}} \det D$$

as well as for the effective theory:

$$\boxed{\text{Tr } e^{-\beta H} = \mathcal{N}_{eff} \int [dU] e^{-\int_0^\beta dx^4 \int d^3x \mathcal{L}_{eff}^{eucl} \{U, v, a, s, p, \theta\}}}$$

Integration extends over all periodic fields:

$$U(\vec{x}, \beta) = U(\vec{x}, 0)$$

- Vertices in the effective Lagrangian remain the same: LEC and CT are independent of T

24.9 Example: quark condensate

- QCD with $N_f \geq 2$ flavours
- ⇒ $\exists N_f^2 - 1$ Nambu-Goldstone-bosons
- Graphs relevant to 3 loops: see Fig. 1 in

Gerber & L. Nucl. Phys. B321 (1989) 387

- Quote the result obtained for massless quarks:

$$\langle \bar{u}u \rangle = \langle 0 | \bar{u}u | 0 \rangle \left\{ 1 - c_1 \frac{T^2}{F_\pi^2} - c_2 \frac{T^4}{F_\pi^4} - c_3 \frac{T^6}{F_\pi^6} \ln \frac{\Lambda_q}{T} + O(T^8) \right\}$$

$$c_1 = \frac{N_f^2 - 1}{12N_f}$$

Binétruy & Gaillard 1985

$$c_2 = \frac{N_f^2 - 1}{288N_f^2}$$

Gasser & L. 1987

$$c_3 = \frac{N_f(N_f^2 - 1)}{1728}$$

Gerber & L. 1989

- If only two quarks are taken massless

$$m_u = m_d = 0 \quad m_s, m_c, \dots \text{ physical}$$

the expansion reads

$$\langle \bar{u}u \rangle = \langle 0 | \bar{u}u | 0 \rangle \left\{ 1 - \frac{T^2}{8F^2} - \frac{T^4}{384F^4} - \frac{T^6}{288F^6} \ln \frac{\Lambda_q}{T} + O(T^8) \right\}$$

- Result is exact: the condensate of massless QCD admits a Taylor series expansion in T . The first few coefficients are determined by F (value of F_π in limit $m_u = m_d = 0$).
- At order T^6 , there is a chiral logarithm; the scale thereof is fixed by the LEC of NLO.

$$\Rightarrow \Lambda_q = 470 \pm 110 \text{ MeV}$$

Gerber & L. 1989

25. Finite volume

25.1 U(1) anomaly

- 3 flavours, singlet axial current:

$$A_\mu = \bar{u}\gamma_\mu\gamma_5 u + \bar{d}\gamma_\mu\gamma_5 d + \bar{s}\gamma_\mu\gamma_5 s$$

- Set $m_u = m_d = m_s = 0$

$$\partial^\mu A_\mu = 6\omega \quad \omega = \frac{1}{16\pi^2} \text{tr}_c G_{\mu\nu} \tilde{G}^{\mu\nu}$$

- \Rightarrow only 8 conserved axial currents
- \Rightarrow only 8 Nambu-Goldstone bosons

- But: ω is a total derivative

$$\partial^\mu A_\mu = \partial^\mu K_\mu \Rightarrow A_\mu - K_\mu \text{ is conserved}$$

This argument looks suspicious because K_μ is not gauge invariant

- Nevertheless, there is a puzzle

Crewther, Christos, Banerjee, ...

25.2 U(1) puzzle

1. For the η' to get a mass, it is essential that the path integral receives contributions from gauge field configurations with nontrivial topology: winding number $\nu = \int dx \omega \neq 0$

2. Configurations with $\nu \neq 0$ give rise to zero modes of the Dirac operator

⇒ For such configurations, $\det D$ tends to zero in the chiral limit, only those with $\nu = 0$ survive

⇒ How can η' retain its mass in chiral limit ???

- To solve the puzzle, need finite volume

L. & Smilga 1992

- Work in euclidean space, hypercube $V = L^4$
- $G_\mu(x)$ periodic up to a gauge transformation, in all four directions

⇒ Winding number quantized: $\nu = 0, \pm 1, \pm 2, \dots$

- Take $N_f = 1$, the problem already arises there
- Turn the external fields off, except for s, p, θ
Take these space-time-independent

- $Z = \int [dG] e^{-\frac{1}{2g^2} \int dx \text{tr} G_{\mu\nu} G^{\mu\nu} + i\theta \int dx \omega} \det D$
 $= Z(g, s, p, \theta, L) \quad g \leftrightarrow \Lambda_{\text{QCD}}$

- Integrate over all configurations with a given ν , then sum over ν

$$Z = \sum_{\nu} e^{i\nu\theta} Z_{\nu}$$

- Z invariant under constant chiral rotations:
see section 8. Ward identities

$$s' + ip' = e^{2i\beta} (s + ip) \quad \theta' = \theta - 2\beta$$

$$Z(g, s', p', \theta', L) = Z(g, s, p, \theta, L)$$

$$\Rightarrow \boxed{Z = Z(g, \sigma, L)} \quad \sigma \equiv (s + ip) e^{i\theta}$$

- If L is large compared to the mass of the lowest state, then the contribution from the ground state dominates:

$$Z = e^{-V\epsilon_0}$$

$$\epsilon_0 = \epsilon_0(g, \sigma) \text{ energy density of ground state}$$

- Parity: $\sigma + \sigma^*$ even, $\sigma - \sigma^*$ odd

\Rightarrow Expansion in powers of σ starts with

$$\epsilon_0(g, \sigma) = \epsilon_{00} + \langle 0 | \bar{q}q | 0 \rangle \text{Re}[\sigma] + O(\sigma^2)$$

Can now turn p off and identify s with the mass of the quark $\Rightarrow \sigma = m e^{i\theta}$

- Result: for $L \gg 1/\Lambda_{\text{QCD}}$, $m \ll \Lambda_{\text{QCD}}$ the partition function of QCD with $N_f = 1$ reads

$$Z = e^{\Sigma V m \cos \theta}$$

where $\Sigma = -\langle 0 | \bar{q}q | 0 \rangle$ is the quark condensate
 V is the volume of the four-dimensional box

25.3 Distribution of winding number

$$e^{\Sigma V m \cos \theta} = \sum_{\nu} e^{i\nu\theta} I_{\nu}(\Sigma V m)$$

$$Z_{\nu} = I_{\nu}(\Sigma V m) \quad \text{Bessel function}$$

- For $\theta = 0$, the partition function is a sum of positive terms, probability for finding winding number ν is

$$p_{\nu} = \frac{Z_{\nu}}{Z} = I_{\nu}(\Sigma V m) e^{-\Sigma V m}$$

- Mean square winding number is $\langle \nu^2 \rangle = \Sigma V m$

$\Sigma V m \ll 1 \rightarrow$ configurations with $\nu = 0$ dominate

$\Sigma V m \gg 1 \rightarrow$ typical winding number is large

- Even for a tiny quark mass, $\Sigma V m$ becomes large if the box is removed
- Topological susceptibility

$$\int dx \langle \omega(x) \omega(0) \rangle = \frac{\langle \nu^2 \rangle}{V} = \Sigma m$$

is independent of the size of the box and $\propto m$

- Extension to $N_f > 1$ and thorough discussion in Phys. Rev. D46 (1992) 5607
- Moral: Suppression of nontrivial topologies is a finite size phenomenon. In the physical situation, where $m_u, m_d, m_s \neq 0$ and $V \simeq \infty$ configurations with $\nu \neq 0$ are not suppressed. Number of zero modes per unit volume is $\propto 1/\sqrt{V}$, hence zero modes are irrelevant.

25.4 Symmetry restoration

- Partition function with $N_f \geq 2$

⇒ ∃ Nambu-Goldstone Bosons

⇒ If the quark masses are zero, the ground state does not dominate the partition function

- In a hypercube of size L , the pions have zero modes: $U(x) = U_0$ constant

Leading contribution to Z for $\theta = 0$:

⇒ $Z = N \int d\mu(U_0) e^{\Sigma V \langle \mathcal{M}(U_0 + U_0^\dagger) \rangle}$

$$\mathcal{M} = \begin{pmatrix} m_u & & \\ & m_d & \\ & & m_s \end{pmatrix}$$

- The relevant parameter is $\Sigma V \mathcal{M}$ also here

- Standard χ PT analysis ("p-regime"):

$$1/L = O(p), \mathcal{M} = O(p^2)$$

$$\Rightarrow \Sigma V \mathcal{M} = O(p^{-2}) \text{ large}$$

\Rightarrow cannot analyze the behaviour in the region where $\Sigma V \mathcal{M}$ is of order 1 in this framework

- For this purpose, we introduced a different bookkeeping: "ε-regime"

$$p = O(\epsilon), 1/L = O(\epsilon), \mathcal{M} = O(\epsilon^4)$$

$$\Rightarrow \Sigma V \mathcal{M} = O(1)$$

- The formula given above represents the leading term in the expansion of the QCD partition function in powers of ϵ :

$$Z = N \int d\mu(U_0) e^{\Sigma V \langle \mathcal{M}(U_0 + U_0^\dagger) \rangle} \{1 + O(\epsilon)\}$$

- Chiral symmetry is restored if $\Sigma V \mathcal{M}$ is small:

$$\mathcal{M} \rightarrow 0 \text{ at finite volume} \Rightarrow \langle 0 | \bar{q}q | 0 \rangle \rightarrow 0$$

- Illustrates general statement: at finite volume, symmetries cannot break down spontaneously

- Symmetry restoration occurs in the ϵ -regime, can study it in detail. Only the NGBs matter if the volume is large enough.
- Method applies to any system which undergoes spontaneous symmetry breakdown.
- Lattice data on the σ -model, for instance, beautifully confirm the predictions.

25.5 Quarks and gluons in a box

- QCD in a 3-dimensional box of size $L \times L \times L$
 $N_f \geq 2$ quark flavours of equal mass m

- Energy levels ?

- Evaluate partition function as a function of T

$$Z = \sum_n e^{-\frac{E_n}{T}}$$

- Energy levels are controlled by $r = F^2 L^3 M$
with $M^2 \equiv 2Bm$ GMOR

⇒ Expand Z in powers of $1/L$, T and m

and order the expansion with

$$1/L = O(\delta), \quad T = O(\delta^3), \quad m = O(\delta^6)$$

⇒ This implies $M = O(\delta^3)$, hence $r = O(1)$

" δ -regime"

- Level spectrum turns out to be the one of a quantum mechanical rotator (freely rotating rigid body)
- Unfortunately, such boxes do not occur in nature, science of the lattice
- This science is very useful and has been developed in various directions. In particular, can study the dependence on the lattice spacing by extending χ PT accordingly

Exercises

1. Evaluate the positive frequency part of the massless propagator

$$\Delta^+(z, 0) = \frac{i}{(2\pi)^3} \int \frac{d^3k}{2k^0} e^{-ikz}, \quad k^0 = |\vec{k}|$$

for $\text{Im } z^0 < 0$. Show that the result can be represented as

$$\Delta^+(z, 0) = \frac{1}{4\pi i z^2}$$

2. Evaluate the d -dimensional propagator

$$\Delta(z, M) = \int \frac{d^d k}{(2\pi)^d} \frac{e^{-ikz}}{M^2 - k^2 - i\epsilon}$$

at the origin and verify the representation

$$\Delta(0, M) = \frac{i}{4\pi} \Gamma\left(1 - \frac{d}{2}\right) \left(\frac{M^2}{4\pi}\right)^{\frac{d}{2}-1}$$

How does this expression behave when $d \rightarrow 4$?

3. Leading order effective Lagrangian:

$$\begin{aligned}\mathcal{L}^{(2)} &= \frac{F^2}{4} \langle D_\mu U D^\mu U^\dagger + \chi U^\dagger + U \chi^\dagger \rangle + h_0 D_\mu \theta D^\mu \theta \\ D_\mu U &= \partial_\mu U - i(v_\mu + a_\mu)U + iU(v_\mu - a_\mu) \\ \chi &= 2B(s + ip) \\ D_\mu \theta &= \partial_\mu \theta + 2\langle a_\mu \rangle \\ \langle X \rangle &= \text{tr} X\end{aligned}$$

- Take the space-time independent part of the external field $s(x)$ to be isospin symmetric (i. e. set $m_u = m_d = m$):

$$s(x) = m \mathbf{1} + \tilde{s}(x)$$

- Expand $U = \exp i\phi/F$ in powers of $\phi = \vec{\phi} \cdot \vec{\tau}$ and check that, in this normalization of the field ϕ , the kinetic part takes the standard form

$$\mathcal{L}^{(2)} = \frac{1}{2} \partial_\mu \vec{\phi} \cdot \partial^\mu \vec{\phi} - \frac{1}{2} M^2 \vec{\phi}^2 + \dots$$

with $M^2 = 2mB$.

- Draw the graphs for all of the interaction vertices containing up to four of the fields $\phi, v_\mu, a_\mu, \tilde{s}, p, \theta$.

4. Show that the classical field theory belonging to the QCD Lagrangian in the presence of external fields is invariant under

$$\begin{aligned}
 v'_\mu + a'_\mu &= V_R(v_\mu + a_\mu)V_R^\dagger - i\partial_\mu V_R V_R^\dagger \\
 v'_\mu - a'_\mu &= V_L(v_\mu - a_\mu)V_L^\dagger - i\partial_\mu V_L V_L^\dagger \\
 s' + ip' &= V_R(s + ip)V_L^\dagger \\
 q'_R &= V_R q_R(x) \\
 q'_L &= V_L q_L
 \end{aligned}$$

where V_R, V_L are space-time dependent elements of $U(3)$.

5. Evaluate the pion mass to NLO of χ PT. Draw the relevant graphs and verify the representation

$$M_\pi^2 = M^2 + \frac{2\ell_3 M^4}{F^2} + \frac{M^2}{2F^2} \frac{1}{i} \Delta(0, M^2) + O(M^6)$$

6. Start from the symmetry property of the effective action,

$$S_{\text{QCD}}\{v', a', s', p', \theta'\} = S_{\text{QCD}}\{v, a, s, p, \theta\} - \int dx \langle \beta \Omega \rangle,$$

and show that this relation in particular implies the Ward identity

$$\begin{aligned}
 \partial_\mu^x \langle 0 | T A_a^\mu(x) P_b(y) | 0 \rangle &= -\frac{1}{4} i \delta(x - y) \langle 0 | \bar{q} \{ \lambda_a, \lambda_b \} q | 0 \rangle \\
 &\quad + \langle 0 | T \bar{q}(x) i \gamma_5 \{ m, \frac{1}{2} \lambda_a \} q(x) P_b(y) | 0 \rangle \\
 a &= 1, \dots, 8, \quad b = 0, \dots, 8
 \end{aligned}$$

7. What is the Ward identity obeyed by the singlet axial current,

$$\partial_\mu^x \langle 0 | T A_0^\mu(x) P_b(y) | 0 \rangle = ?$$