

**X-RAY CRYSTALLOGRAPHIC
STUDIES OF A FEW ADRENERGIC
DRUGS AND THEIR INTERACTIONS**

Thesis submitted to the Jadavpur University in partial
fulfilment for the degree of
DOCTOR OF PHILOSOPHY (Science)
by Rakhi Sengupta

Rakhi Sengupta, M.Sc.
Senior Research Fellow,
Crystallography and Molecular Biology Division,
Saha Institute of Nuclear Physics,
1/AF Bidhannagar, Calcutta - 700064
India

CONTENTS

Preface

Part I

Chapter

1	Adrenergic drugs: a brief overview	
1.1	Introduction	1
1.2	Subtype selective adrenergic drugs	5
1.3	Therapeutic applications	9
1.4	Structure-activity studies	10
1.4.1	Phenethylamines	11
1.4.2	Imidazolines	20
2	Crystal structure of an adrenergic agonist: Mephentermine hemisulfate monohydrate	
2.1	Introduction	33
2.2	Experimental	33
2.3	Structure solution and refinement	37
2.4	Results and discussion	
2.4.1	Bond lengths and angles	43

2.4.2	Conformation	48
2.4.3	Hydrogen bonding and molecular packing	49
3	Crystal structure of an α -adrenergic agonist: Oxymetazoline hydrochloride monohydrate	
3.1	Introduction	51
3.2	Experimental	51
3.3	Structure solution and refinement	55
3.4	Results and discussion	
3.4.1	Bond lengths and angles	60
3.4.2	Conformation	63
3.4.3	Mean plane	65
3.4.4	Hydrogen bonding and molecular packing	66
4	Crystal structure of β -adrenergic agonists: Terbutaline hemisulfate and methoxyphenamine hydrochloride	
4.1	Introduction	67
4.2	Experimental	
	(a) Terbutaline hemisulfate	67
	(b) Methoxyphenamine hydrochloride	76
4.3	Results and discussion	
4.3.1	Bond lengths and angles	83
4.3.2	Conformation	91
4.3.3	Hydrogen bonding and molecular packing	92

5	Summary and conclusion	
5.1	Phenethylamines	
	a) Structural and conformational features	95
	b) Crystal packing and molecular interactions	100
5.2	Imidazolines	
	a) Structural and conformational features	101
	b) Crystal packing and molecular interactions	106
5.3	Comparison between phenethylamines and imidazolines	107
5.4	Conclusion	108

Part II

Chapter

6	G-protein coupled receptors (GPCRs)	
6.1	Introduction	110
6.2	GPCR linked signal transduction pathway	111
6.3	Structure of G proteins	111
6.4	Different types of G protein coupled receptors	113
6.4.1	Adrenergic receptors	113
6.5	Biochemical studies of GPCRs	115
6.6	Ligand binding and G protein activation regions	116
6.7	Structure of GPCRs	118
6.8	Structure of Bacteriorhodopsin	118
6.9	Structure of Rhodopsin	120

6.10	Modeling of GPCRs	120
6.11	Purpose of 3D modeling of α_{2A} - adrenergic receptor from human platelets	123
7	Modeling of α_{2A} - adrenergic receptor from human platelets and study of ligand- receptor interactions	
7.1	Introduction	124
7.2	Identification of the seven TM helical regions	
	a) Hydrophobicity plots	125
	b) Algorithm of Eisenberg, Schwarz, Komaromy and Wall	126
	c) Method of Klein, Kanehisa and DeLisi	127
7.2.1	Results and discussion	128
7.3	Helix construction and packing	
7.3.1	Principles of helix packing	129
7.3.2	Surface complementarity function	130
7.3.3	S_c calculations for bacteriorhodopsin	131
7.3.4	Helix topology	132
7.3.5	Helix construction	133
7.3.6	Helix packing to form hepta-helical bundle	
	a) Coarse grid conformational search	134
	b) Construction of the helix-bundle	137
	c) Fine grid conformational search	137
	d) Construction of loops	138
7.4	Discussion of the model	138
7.5	Docking of ligands to α_{2A} - AR model	
7.5.1	Site-directed mutagenesis data of α_{2A} - AR	140

7.5.2	Ligand docking	143
7.5.3	Results and discussion	
	a) Docking of phenethylamines	144
	b) Docking of imidazolines	145
	c) Important interhelical contacts	145
	d) Attempts to dock a few β specific drugs	146
7.6	Conclusion	146
	<i>References</i>	