

Study of Charge Density Wave Transition in Electron-Phonon Interaction Models

Sanjoy Datta

Theoretical Condensed Matter Physics Division,
Saha Institute of Nuclear Physics, 1/AF, Bidhannagar, Kolkata 700064

Email: sanjoy.datta@saha.ac.in



Thesis submitted for the Degree of
Doctor of Philosophy in Science (Physics)
Jadavpur University, Jadavpur, Kolkata, India
2009

Contents

Acknowledgments	vi
List of Publications	ix
Preface	x
1 INTRODUCTION	1
1.1 Peierls Transition	2
1.2 Charge Density Wave Physics	5
1.3 One-dimensional CDW systems	12
1.4 Higher-dimensional CDW systems	14
1.5 Coexistence of CDW and superconductivity	14
1.5.1 Supersolidity	15
1.6 Luttinger Liquid	17
1.7 Bosonization	19
1.8 Anisotropic Heisenberg Model	23
1.9 Modified Lanczos Method	24
Bibliography	28
2 Electron-Phonon Interaction Models	29
2.1 Fröhlich Hamiltonian	30
2.2 Holstein Hamiltonian	33
2.2.1 Lang-Firsov Transformation	34
2.2.2 Charge density wave transition in the Holstein Hamiltonian	36
Bibliography	38

3	Peierls instability in the SOHM in the anti-adiabatic regime	39
3.1	Effective polaronic Hamiltonian	40
3.2	Luttinger liquid to CDW transition	43
3.3	Results and discussions	44
	Bibliography	53
4	Peierls instability in the SOHM in the adiabatic regime	54
4.1	Phonon softening in the LL phase	55
4.2	Half-filled case	57
4.3	Less than half-filling	61
4.4	T=0 phase diagram at non-half-filling	63
	Bibliography	67
5	Quantum phase transition in Bose-Holstein model	68
5.1	Effective Hamiltonian	70
5.2	Mean field analysis	71
5.3	Diagonal long range order and structure factor	76
5.3.1	$J_2 = 0$	77
5.3.2	$J_2 \neq 0$	79
5.4	Off-diagonal long range order	81
5.4.1	Condensate fraction	82
5.4.2	Superfluid fraction	83
5.5	Energy consideration	84
5.5.1	$J_2 = 0$	84
5.5.2	$J_2 \neq 0$	84
5.6	Results and discussions	86
	Bibliography	97
6	Conclusions and future prospects	98
		100
A	Lang-Firsov Transformation	101

B	103
B.1	103
B.2	104
B.2.1	The set of degenerate states for half-filled case 104
B.2.2	Derivation of energy E_{Φ_n} in the half-filled case 105
B.3	106
B.3.1	The case when $\epsilon_{\vec{k}-\vec{p}} - \epsilon_{\vec{k}} = \epsilon_{\vec{k}-2\vec{p}} - \epsilon_{\vec{k}-\vec{p}} = \omega_0$ 109
B.3.2	When both the solutions exist 109