

Role of Huntingtin (Htt) interacting proteins in apoptosis induction in cultured cells expressing exon1 of mutated Huntingtin gene

**THESIS SUBMITTED FOR THE DEGREE OF DOCTOR OF PHILOSOPHY
(SCIENCE) OF JADAVPUR UNIVERSITY, KOLKATA, INDIA**



Manisha Banerjee (nee Bhattacharya)

CRYSTALLOGRAPHY & MOLECULAR BIOLOGY DIVISION

SAHA INSTITUTE OF NUCLEAR PHYSICS

KOLKATA, INDIA

2009

Content

Abbreviations	1
Synopsis	4
Chapter 1: General Introduction	
1.01 Huntington's disease	15
1.02 Genetic Basis of the Disease: mutation to the gene <i>Huntingtin (HTT)</i>	15
1.03 Domain organization of Huntingtin protein: insight to its function	16
1.04 Molecular pathogenesis of Huntington's Disease	17
1.04.01 Aggregates of mutant HTT proteins	19
1.04.02 Apoptosis	20
1.04.03 Mitochondrial dysfunction	21
1.04.04 Autophagy and Ubiquitin proteosomal degradation system in HD	21
1.04.05 Transcriptional deregulation in HD	22
1.04.06 Possible mechanisms of transcription deregulation	26
1.05 Huntingtin (HTT) Interacting Proteins	30
1.06 Role of HTT-interacting proteins in alteration of mutant HTT induced aggregates and toxicity	33
1.07 Sequestered poly Q Hypothesis	36
1.08 Huntingtin interacting protein 1 (HIP-1) and its molecular partner HIPPI	
1.08.01 Structural feature of Huntingtin interacting protein 1 (HIP-1): Clue to its function	37
1.08.02 HIP-1 interacting proteins: participation in apoptosis and survival	38
1.08.03 HIP-1: role in protein trafficking and transcription	41
1.08.04 Structural organization of HIPPI, its interacting partners, and involvement in apoptosis	41
1.08.05 Role of HIPPI in transcription regulation	43
1.09 Myeloid Leukemia Factors (MLFs): its interacting partners and possible functions	44

Chapter 2: Materials and Methods

2.01 Chemicals	47
2.02 Mammalian cells, Bacterial strains and Plasmids	
2.02.01 Mammalian Cells	48
2.02.02 Bacterial Strains	48
2.02.03 Plasmids	48
2.02.04 Human Brain plasmid cDNA library	51
2.03 Methods	
2.03.01 Mammalian cell culture	51
2.03.02 Sterilization	51
2.03.03 Transfection of mammalian cells	51
2.03.04 RNA Isolation and first strand coding DNA (cDNA) preparation	52
2.03.05 Gene cloning	53
2.03.06 Site directed mutagenesis	58
2.03.07 PCR product cloning in pTZ57R/T vector	58
2.03.08 Different microscopic methods	59
2.03.09 Detection of Apoptosis	61
2.03.10 Gene expression study by Real Time PCR	62
2.03.11 Western blot analysis	63
2.03.12 Immunoprecipitation assay	64
2.03.13 <i>In vivo</i> chaperone activity assay	66
2.03.14 Protein expression, extraction and purification	67
2.03.15 Circular Dichroism (CD) spectral analysis	68
2.03.16 Crystallographic technique	69
2.03.17 Modeling	69
2.03.18 Procedures for Electrophoresis	70
2.03.19 Filter retardation assay	70
2.03.20 Bioinformatics approaches to analyze HTT interacting proteins	71

Chapter 3: Huntingtin interacting proteins: involvement in diverse molecular functions, biological processes and pathways

3.01 Expressions of HTT-interacting proteins in brain	77
3.02 Influence of poly Q length at HTT for the interactions with other proteins	78
3.03 HTT-interacting proteins: their molecular functions	81
3.04 HTT-interacting proteins: their involvement in biological processes	83
3.05 HTT-interacting proteins involve in diverse biological pathways	85
3.06 Involvement of HTT-interacting proteins in other diseases	89

Chapter 4: Modulation of mutated N-terminal HTT mediated aggregates and apoptosis by HTT interacting proteins

PART 4A: Supression of mutant N-terminal HTT aggregates, cellular toxicity and apoptosis by human myloid leukemia factors MLF1 and MLF2

4A.01 Preferential interaction of Human myloid leukemia factors hMLF1 and hMLF2 with mutated N-terminal HTT protein coded by exon1 of <i>HTT</i> gene.	
4A.01.01 Expression and localization of MLF1 and MLF2 with the N-termial HTT coded by <i>HTT</i> exon1.	107
4A.01.02 Interaction of MLF1 and MLF2 with N-terminal HTT coded by <i>HTT</i> exon-1 by immunoprecipitation.	108
4A.02 Modulation of aggregates formed by N-terminal HTT with 83Q by co-expression of GFP-MLF1 and GFP-MLF2	
4A.02.01 Decreased number of cells containing mutated HTT aggregates in the presence of GFP-MLF1 or GFP-MLF2	110
4A.02.02 Live cell microscopy reveals co-expression of GFP-MLF1 reduces poly Q mediated aggregation kinetics	110
4A.02.03 Reduction of SDS insoluble aggregates in presence of MLF1 and MLF2	111
4A.02.04 Fluorescence recovery after photo bleaching (FRAP) of HTT aggregates in the presence of either GFP-MLF1 or GFP-MLF2	112

4A.03 Reduction of DsRed-83Q induced apoptosis by co-expression of GFP-MLF1 and GFP-MLF2	114
4A.04 Partial reduction of mutated HTT mediated aggregation and apoptosis in presence N-terminal MLF1	
4A.04.01 Generation of deletion mutants of MLF1	116
4A.04.02 Expression of deletion mutants of MLF1 separately and together with HTT exon 1 in Neuro-2A cell	117
4A.04.03 Interaction of N-terminal MLF1 with N-terminal HTT coded by the exon1 of HTT gene	118
4A.04.04 Reduction of mutated N-terminal HTT aggregates by domain “A” (5-100aa) of MLF1	119
4A.04.05 Reduction of toxicity and apoptosis in DsRed-83Q expressing cells by domain “A” (5-100aa) of MLF1	119
4A.04.06 Fluorescence recovery after photo bleaching (FRAP) of Htt aggregates in the presence and absence of GFP-“A” and GFP-“B”.	120
4A.05 Association of transcription regulators CREB, CBP and p53 with mutated HTT aggregates and release of these transcription regulators in the presence of GFP-MLF1	
4A.05.01 Release of CREB, CBP and p53 from HTT aggregates by MLF1	121
4A.05.02 Reversal of expression of GADD45A in the presence of GFP-MLF1	122
4A.05.03 In vivo chaperone activity of MLF1	123

PART 4B: Role of DNAJ (Hsp40) homolog, subfamily B, member 3 (DNAJB3, also known as HCG3) in the reduction of mutated N-terminal HTT coded by *HTT* gene mediated aggregation and apoptosis

4B.01 Theoretical analysis of DNAJB3 sequence	
4B.01.01 Comparison with its homologous chaperone	130
4B.01.02 Structural organization of different domains of DNAJB3	130
4B.02 Co-localization of HCG3 with wild type as well as mutant N-terminal HTT protein with 83Q coded by exon1 of HTT gene	

4B.02.01 Colocalization of HCG3 with N-terminal HTT coded by exon-1 having 83Q or 16Q.	132
4B.02.02 Physical interaction of HCG3 with HTT exon1 containing 16Q and 83Q	132
4B.03 Expression of GFP-HCG3 reduces the mutant N-terminal HTT induced aggregates	
4B.03.01 Suppression of aggregates formed by N-terminal HTT with 83Q by co-expression of GFP-HCG3	134
4B.03.02 Reduction of SDS insoluble aggregates formed by DsRed-83Q in presence of GFP-HCG3	134
4B.04 Reduction of mutated <i>HTT</i> exon 1 induced apoptosis by GFP-HCG3	135
4B.05 HCG3 is an active component of chaperone machinery	
4B.05.01 Co localization of HCG3 with Hsp70	136
4B.05.02 <i>In vivo</i> chaperone activity of HCG3	137

PART 4C: Role of Huntingtin Interacting Protein 1(HIP-1) in the modulation of aggregates and apoptosis in cells expressing mutated N-terminal HTT coded by the *HTT* exon1

4C.01 Reduction of mutated N-terminal HTT coded by the exon1 of HTT gene mediated aggregates formation and apoptosis induction by HIP-1	
4C.01.01 Expression of HIP-1 and its N terminal domain separately and along with <i>HTT</i> exon1	142
4C.01.02 Suppression of expanded N-terminal HTT mediated aggregation formation by HIP-1	144
4C.01.03 Reduction of mutated N-terminal HTT mediated apoptosis by HIP-1	145

Chapter 5: Transcription regulation by HIP-1 and HIPPI

5.01 Structural analysis of pseudo Death Effector Domain (pDED) of HIPPI	
5.01.01 Construction of recombinant pDED-HIPPI	151
5.01.02 Determination of molecular weight of pDED-HIPPI by mass spectrometry	151
5.01.03 CD spectroscopy of 6NH tagged pDED-HIPPI	151

5.01.04 Crystallization of pDED-HIPPI	152
5.01.05 X-ray Data collection and processing	154
5.01.06 Preliminary crystallographic analysis of pDED-HIPPI	155
5.01.07 Modeling of pDED of HIPPI: possible residues involved in the interaction with DNA	156
5.01.08 PreDs rule prediction for DNA binding domain of pDED-HIPPI	157
5.02 Involvement of 393 arginine of HIPPI in the transcriptional regulation of Caspase1 and apoptosis	
5.02.01 Exogenous expression of wild type (R393) and mutant (E393) pDED of HIPPI in HeLa cell	158
5.02.02 Physical interaction of HIP-1 with the wild type and mutant pDED and recruitment of pro-caspase-8	158
5.02.03 Reduction of caspase-1 gene expression in cells expressing mutated pDED	159
5.02.04 Apoptosis induction by mutant pDED	159
5.02.05 Mutant pDED-HIPPI (R393E) can not interact with caspase-1 promoter	159
5.03 Role of HIP-1 in the transcriptional regulation of caspase-1 gene by translocating HIPPI into nucleus	
5.03.01 Expression of caspase-1 in HIP-1 reduced HIPPI expressing cell	161
5.03.02 Reduction of Apoptosis induction in HIP-1 reduced HIPPI expressing cell	163
5.03.03 Translocation of HIPPI into the nucleus requires expression of HIP-1 in HeLa	164
5.03.04 Translocation of the pDED and its mutant to the nucleus by HIP-1	166
5.03.05 Endogenous HIPPI interacts with caspase-1 promoter	168
5.03.06 Endogenous HIPPI increases the expression level of caspase-1	168
5.03.07 Translocation of endogenous HIPPI into nucleus requires HIP-1 in N2a	169
5.03.08 Exogenous over expression of HIP-1 increases caspase-1 gene expression in N2a cell	170

CHAPTER 6: Summary and conclusion	176
Bibliography	183
Appendix	i-xi